

Product Information: Glucerna® 1.2 Cal

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Glucerna® 1.2 Cal

Specialized Nutrition for Enhanced Glycemic Control

- For patients with type 1 or type 2 diabetes.
- For patients with hyperglycemia/abnormal glucose tolerance resulting from metabolic stress, such as illness, trauma or infection.
- For tube or oral feeding.
- For supplemental or sole-source nutrition.
- Use under medical supervision.
- GLUCERNA 1.2 CAL has been clinically shown to significantly reduce peak postprandial blood glucose response by 49 mg/dL (69%) and reduce glycemic variability versus a standard 1.2-Cal formulation in people with diabetes.^{1,2}
- Caloric distribution - protein, 20%; fat, 45%; carbohydrate, 35%.
- 20% of calories from protein to promote anabolism and support wound healing.
- At 1.2 Cal/mL, patient needs can be met with less volume.
- Meets or exceeds 100% of the RDIs for 24 key vitamins and minerals in 1500 Cal (1250 mL).
- 4% of total calories from saturated fatty acids and rich in monounsaturated fatty acids (28% of total calories), consistent with the American Heart Association and American Diabetes Association guidelines.³
- Provides plant-based omega-3 fatty acids from canola oil (3 g of ALA per 1500 Cal) and the omega-3 fatty acids EPA and DHA (1 g per 1500 Cal) from fish oil consistent with American Heart Association reported levels to support cardiovascular health.⁴
- Slowly digested carbohydrate sources for enhanced glycemic response.
- Has chromium picolinate to help support glucose levels.
- Chromium from chromium picolinate to help the body's insulin work.
- Kosher.
- Halal.
- Gluten-free; suitable for lactose intolerance.



Safety Precautions

- Not for patients with galactosemia.
- Not for IV use.
- **Feeding sets are for single-patient use only. Use clean technique to avoid set and or product contamination**

¹ Data on File: Study BK06 Part B. Abbott Laboratories, 2008.

² Alish C, et al. *Diabetes Technol Ther* 2010;12:419.

³ Buse JB, et al. *Circulation* 2007;115:114-126.

⁴ Kris-Etherton PM, et al. *Circulation* 2002;106:2747-57.

Ingredients

Vanilla Ready to Drink:

Water, Sodium Caseinate, Corn Maltodextrin, High Oleic Safflower Oil, Isomaltulose, Canola Oil, Fructose, Soy Protein Isolate, Sucromalt, Short-Chain Fructooligosaccharides, Glycerine, Milk Protein Concentrate, Oat Fiber, Soy Lecithin, Soy Fiber, Potassium Citrate, Marine Oil (May Contain One or More of the Following: Anchovy, Menhaden, Salmon, Sardine, Tuna), Magnesium Phosphate, Natural & Artificial Flavors, Potassium Chloride, m-Inositol, Calcium Carbonate, Calcium Citrate, Sodium Citrate, Ascorbic Acid, Choline Chloride, Sodium Chloride, L-Carnitine, Taurine, Carrageenan, Ferrous Sulfate, dl-Alpha-Tocopheryl Acetate, Zinc Sulfate, Niacinamide, Calcium Pantothenate, Manganese Sulfate, Cupric Sulfate, Vitamin A Palmitate, Thiamine Chloride Hydrochloride, Pyridoxine Hydrochloride, Beta-Carotene, Riboflavin, Chromium Picolinate, Folic Acid, Biotin, Sodium Molybdate, Sodium Selenate, Potassium Iodide, Phylloquinone, Cyanocobalamin, and Vitamin D3.

Allergens: Contains milk and soy ingredients.

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Glucerna® 1.2 Cal

Specialized Nutrition for Enhanced Glycemic Control

Availability

List Number	Item
50902	Glucerna 1.2 Cal Ready-To-Hang Institutional / 1500-mL (50.7-fl-oz) Bottle / 6 ct (discontinued)
50904	Glucerna 1.2 Cal Ready-to-Drink Vanilla Institutional / 8-fl-oz (237-mL) Can / 24 ct
50906	Glucerna 1.2 Cal Ready-To-Hang Institutional / 1000-mL (33.8-fl-oz) Bottle / 8 ct (discontinued)
62352	Glucerna 1.2 Cal Non-Spikable Cap Ready-To-Hang Institutional / 1500-mL (1.6-QT) Bottle / 6 ct
62675	Glucerna 1.2 Cal Cross Cap Ready-To-Hang Institutional / 1500-mL (1.6-QT) Bottle / 6 ct
62677	Glucerna 1.2 Cal Cross Cap Ready-To-Hang Institutional / 1000-mL (1.1-QT) Bottle / 8 ct

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Glucerna® 1.2 Cal

Specialized Nutrition for Enhanced Glycemic Control

Nutrition Information - Vanilla Ready to Drink

	8 fl oz (237 mL)	
	Value	%RDI* / %DV
Nutrient Density, Cal/mL	1.2	
Protein, % Cal	20.0	
Fat, % Cal	45.0	
Monounsaturated Fatty Acids, % Cal	28.0	
Carbohydrate, % Cal	35.0	
Total Cal:g Nitrogen	125:1	
Nonprotein Cal:g Nitrogen	100:1	
Osmolality, mOsm/kg H ₂ O	720	
Renal Solute Load, mOsm/L	479	
Viscosity	Thin (room temperature, chilled)	
Minimum Tube Size for Gravity/Pump Feeding (Fr)	10/8	
Exchanges per 8 fl oz [†]	2 Starch, 2 Medium-Fat Meat OR 2 Starch, 1 Medium-Fat Meat, 1 Fat	
Carb Choices	2	
Protein, g	14.2	
Fat, g	14.2	
Carbohydrate, g	27.1	
Dietary Fiber, g	3.8 [‡]	
Taurine, mg	32	
L-Carnitine, mg	42	
m-Inositol, mg	200	
Water, g	192	
Calories	285	
Vitamin A, IU	1840 [§]	37
Vitamin D, IU	82	21
Vitamin E, IU	9.2	31
Vitamin K, mcg	24	30
Vitamin C, mg	62	105
Folic Acid, mcg	76	19
Vitamin B1, mg	0.29	19
Vitamin B2, mg	0.33	19
Vitamin B6, mg	0.38	19
Vitamin B12, mcg	1.2	20
Niacin, mg	3.8	19
Choline, mg	105	Not established
Biotin, mcg	57	19
Pantothenic Acid, mg	1.9	19
Beta-Carotene, mg	0.63	Not established
Sodium, mg	265	Not established

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Nutrition Information - Vanilla Ready to Drink

	8 fl oz (237 mL)	
	Value	%RDI* / %DV
Potassium, mg	480	Not established
Chloride, mg	305	9
Calcium, mg	190	19
Phosphorus, mg	190	19
Magnesium, mg	76	19
Iodine, mcg	29	19
Manganese, mg	0.38	19
Copper, mg	0.38	19
Zinc, mg	2.9	19
Iron, mg	3.5	19
Selenium, mcg	14	20
Chromium, mcg	38	32
Molybdenum, mcg	19	25

Vanilla Ready to Drink Footnotes & References

Per 8 fl oz (237 mL)

*RDI percentages for adults and children 4 or more years of age.

†Exchange Lists for Diabetes. Alexandria, Va, and Chicago: American Diabetes Association and American Dietetic Association, 2008.

‡1.4 g of total dietary fiber from soy, oat and corn fibers; 2.4 g of short-chain fructooligosaccharides.

§830 IU/8 fl oz of vitamin A activity supplied by 0.63 mg of beta-carotene.

Preparation

Instructions for Use: 8-fl-oz can

- Store unopened at room temperature; avoid extreme temperatures.
- Shake well. Glucerna 1.2 is ready to use.
- Clean top of container thoroughly before opening.
- Once opened, cover, refrigerate and **use within 48 hours**.

Oral Feeding

- May be used for total or supplemental nutrition.
- May be fed at room temperature or chilled.

Tube Feeding

- Follow physician's instructions.
- Adjust flow rate and volume according to patient's condition and tolerance.
- Gravity feed at room temperature with a 10 Fr or larger tube; use a pump set with smaller tubes.
- Additional fluid requirements should be met by giving water between or after feedings, or when flushing the tube.
- Avoid contamination during preparation and use.

Instructions for Use: Ready-To-Hang Container

All liquid medical foods, regardless of type of administration system, require careful handling because they can support microbial growth. Follow these instructions for clean technique and proper setup to reduce the potential for microbial contamination. NOTE: Failure to follow the **Instructions for Use** increases the potential for microbial contamination and reduces hangtime.

- Pump feeding is recommended; use an 8 Fr or larger tube. If gravity feeding use a 10 Fr or larger tube.
- Administer product at room temperature.
- THOROUGHLY wash hands with soap and water before handling container or feeding set.
- Turn container upside down and SHAKE VIGOROUSLY, using a twisting motion for at least 10 seconds.
- DO NOT touch any part of the container or feeding set that comes into contact with the formula.
- When initiating feeding, follow physician's instructions. Adjust flow rate and volume according to patient's condition and tolerance.
- Additional fluid requirements should be met by giving water between or after feedings or when flushing the tube.
- Follow Directions for Use provided with Abbott feeding set.

For Use with Abbott Piercing Pin Set:

- Lift tab to remove sticker from cap, taking care NOT to touch port. DO NOT remove the plastic cap.
- Insert spike into preattached cap, completely pierce foil, and turn 1/4 turn.
- Close clamp on set before inverting container.
- Invert container and suspend, using the hanging ring on the bottom of the container.
- **Hang product up to 48 hours after initial spike when clean technique and only one new set is used. Otherwise, hang no longer than 24 hours.**

For Use with Abbott Safety Screw Connector Set:

- Remove the dust cover from the RTH Safety Screw Cap.
- Remove the dust cover from the Safety Screw Connector Set.
- Insert the Safety Screw Connector Set into the port on the RTH Safety Screw Cap, completely pierce foil.
- Turn the Safety Screw Connector Clockwise until it is securely fastened.
- Close clamp on set before inverting container.
- Invert container and suspend, using the hanging ring on the bottom of the container.
- **Hang product up to 48 hours when clean technique and only one Screw Cap feeding set is used.**

For Use with Abbott Screw Cap Set:

- Remove plastic cap from RTH container taking care NOT to touch the foil seal.
- Visually inspect foil for signs of leakage. DO NOT use if leakage is apparent.
- Secure 40 mm Screw Cap set onto the RTH container.
- Close clamp on set before inverting container.
- Invert container and suspend, using the hanging ring on the bottom of the container.
- **Hang product up to 24 hours when clean technique and only one Screw Cap feeding set is used.**