

Product Information: Jevity® 1.5 Cal

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Jevity® 1.5 Cal

High-Protein, High-Calorie Nutrition With Fiber and NutraFlora® scFOS®

- JEVITY 1.5 CAL is a calorically dense formula with a unique fiber blend that provides Complete, Balanced Nutrition®.
- For tube feeding.
- For supplemental or sole-source nutrition.
- May be used for oral feeding of patients with altered taste perception.
- Use under medical supervision.
- Concentrated calories (1.5 Cal/mL) to help tube-fed patients gain and maintain healthy weight.
- 22 grams of dietary fiber in 1 Liter (12 g of a unique soluble and insoluble fiber blend and 10 g of short-chain fructooligosaccharides [scFOS]), which helps meet daily fiber needs. Fiber helps moderate bowel function.
- 15 g NutraFlora® scFOS®/1500 mL. scFOS are prebiotics that stimulate the growth of beneficial bacteria in the colon.
- Meets or exceeds 100% of the RDI for 24 essential vitamins and minerals in 1500 Cal (1 Liter).
- Halal.
- Kosher.
- Gluten-free; suitable for lactose intolerance.



Safety Precautions

- Not for patients with galactosemia.
- Not for IV use.
- **Feeding sets are for single-patient use only. Use clean technique to avoid set and/or product contamination**

* NutraFlora® and scFOS® are not registered trademarks of Abbott Laboratories.

Ingredients

Unflavored:

Water, Corn Maltodextrin, Corn Syrup Solids, Sodium & Calcium Caseinates, Canola Oil, Corn Oil, Soy Protein Isolate, Fructooligosaccharides, Medium-Chain Triglycerides, Oat Fiber, Soy Fiber, Potassium Citrate, Calcium Phosphate, Gum Arabic, Sodium Citrate, Magnesium Phosphate, Soy Lecithin, Magnesium Chloride, Carboxymethylcellulose, Ascorbic Acid, Choline Chloride, Potassium Chloride, Taurine, L-Carnitine, Zinc Sulfate, dl-Alpha-Tocopheryl Acetate, Ferrous Sulfate, Niacinamide, Calcium Pantothenate, Manganese Sulfate, Cupric Sulfate, Thiamine Chloride Hydrochloride, Pyridoxine Hydrochloride, Riboflavin, Vitamin A Palmitate, Folic Acid, Biotin, Chromium Chloride, Sodium Molybdate, Potassium Iodide, Sodium Selenate, Phylloquinone, Cyanocobalamin, and Vitamin D3.

Allergens: Contains milk and soy ingredients.

Availability

List Number	Item
57329	Jevity 1.5 Cal Ready-To-Hang Institutional / 1000-mL (33.8-fl-oz) Bottle / 8 ct (discontinued)
57331	Jevity 1.5 Cal Ready-To-Hang Institutional / 1500-mL (50.7-fl-oz) Bottle / 6 ct (discontinued)
57333	Jevity 1.5 Cal Ready-to-Drink Institutional / 8-fl-oz (237-mL) Can / 24 ct
62681	Jevity 1.5 Cal Cross Cap Ready-To-Hang Institutional / 1000-mL (1.1-QT) Bottle / 8 ct
62689	Jevity 1.5 Cal Cross Cap Ready-To-Hang Institutional / 1500-mL (1.6-QT) Bottle / 6 ct

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Nutrition Information - Unflavored

	8 fl oz (237 mL)		1000 mL		1500 mL	
	Value	%RDI* / % DV	Value	%RDI* / % DV	Value	%RDI* / % DV
Nutrient Density, Cal/mL	1.5		1.5		1.5	
Protein, % Cal	17.0		17.0		17.0	
Carbohydrate, % Cal	53.6		53.6		53.6	
Fat, % Cal	29.4		29.4		29.4	
MCT:LCT	19:81		19:81		19:81	
Cal to Meet 100% RDIs	1500		1500		1500	
mL to Meet 100% RDIs	1000		1000		1000	
Total Cal:g Nitrogen	147:1		147:1		147:1	
Nonprotein Cal:g Nitrogen	122:1		122:1		122:1	
Osmolality, mOsm/kg H ₂ O	525		525		525	
Renal Solute Load, mOsm/L	518		518		518	
Viscosity	Nectar-Like (Room Temperature and Chilled)		Nectar-Like (Room Temperature and Chilled)		Nectar-Like (Room Temperature and Chilled)	
Minimum Tube Size for Gravity/Pump Feeding (Fr)	12/10		12/10		12/10	
Protein, g	15.1		63.8		95.7	
Fat, g	11.8		49.8		74.7	
Carbohydrate, g	51.1		215.7		323.6	
Dietary Fiber, g	5.3 [†]		22.0 [†]		33.0 [†]	
L-Carnitine, mg	36		150		225	
Taurine, mg	36		150		225	
Water, g/mL/cc	180		760		1140	
Calories	355		1500		2250	
Vitamin A, IU	1185	24	5000	100	7500	150
Vitamin D, IU	95	24	400	100	600	150
Vitamin E, IU	11	37	45	150	68	225
Vitamin K, mcg	19	24	80	100	120	150
Vitamin C, mg	72	120	300	500	450	750
Folic Acid, mcg	145	36	600	150	900	225
Thiamin (Vitamin B1), mg	0.54	36	2.3	155	3.4	225
Riboflavin (Vitamin B2), mg	0.61	36	2.6	155	3.9	230
Vitamin B6, mg	0.72	36	3.0	150	4.5	225
Vitamin B12, mcg	2.2	37	9.0	150	14	235
Niacin, mg	7.2	36	30	150	45	225
Choline, mg	145	Not established	600	Not established	900	Not established
Biotin, mcg	110	37	450	150	675	225
Pantothenic Acid, mg	3.6	37	15	150	23	230

For more information, contact your Abbott Nutrition Representative or visit www.abbottnutrition.com

Nutrition Information - Unflavored

	8 fl oz (237 mL)		1000 mL		1500 mL	
	Value	%RDI* / % DV	Value	%RDI* / % DV	Value	%RDI* / % DV
Sodium, mg	330	Not established	1400	Not established	2100	Not established
Sodium, mEq	14.3		60.9		91.3	
Potassium, mg	510	Not established	2150	Not established	3225	Not established
Potassium, mEq	13.1		55.1		82.7	
Chloride, mg	320	9	1360	40	2040	60
Chloride, mEq	9.0		38.4	120	57.6	
Calcium, mg	285	29	1200	120	1800	180
Phosphorus, mg	285	29	1200	100	1800	180
Magnesium, mg	95	24	400	100	600	150
Iodine, mcg	36	24	150	100	225	150
Manganese, mg	1.2	60	5.0	250	7.5	375
Copper, mg	0.48	24	2.0	100	3.0	150
Zinc, mg	5.4	36	23	155	34	225
Iron, mg	4.3	24	18	100	27	150
Selenium, mcg	17	24	70	100	105	150
Chromium, mcg	29	24	120	100	180	150
Molybdenum, mcg	36	48	150	200	225	300

Unflavored Footnotes & References

Per 8 fl oz (237 mL)

*RDI percentages for adults and children 4 or more years of age.

†2.9 g of total dietary fiber from a unique fiber blend and 2.4 g of fructooligosaccharides.

Per 1000 mL

*RDI percentages for adults and children 4 or more years of age.

†12.0 g of total dietary fiber from a unique fiber blend and 10.0 g of fructooligosaccharides.

Per 1500 mL

*RDI percentages for adults and children 4 or more years of age.

†18.0 g of total dietary fiber from a unique fiber blend and 15.0 g of fructooligosaccharides.

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Abbott Nutrition Abbott Laboratories
 Columbus, OH 43219-3034
 1-800-227-5767



Preparation

Instructions for Use: Ready-To-Hang Container

All liquid medical foods, regardless of type of administration system, require careful handling because they can support microbial growth. Follow these instructions for clean technique and proper setup to reduce the potential for microbial contamination. NOTE: Failure to follow the **Instructions for Use** increases the potential for microbial contamination and reduces hangtime.

- Pump feeding is recommended; use a 10 Fr or larger tube. If gravity feeding, use a 12 Fr or larger tube.
- Administer product at room temperature.
- THOROUGHLY wash hands with soap and water before handling container or feeding set.
- Turn container upside down and SHAKE VIGOROUSLY, using a twisting motion for at least 10 seconds.
- DO NOT touch any part of the container or feeding set that comes into contact with the formula.
- When initiating feeding, follow physician's instructions. Adjust flow rate and volume according to patient's condition and tolerance.
- Additional fluid requirements should be met by giving water between or after feedings or when flushing the tube.
- Follow Directions for Use provided with Abbott feeding set.

For Use with Abbott Safety Screw Connector Set:

- Remove the dust cover from the RTH Safety Screw Cap.
- Remove the dust cover from the Safety Screw Connector Set.
- Insert the Safety Screw Connector Set into the port on the RTH Safety Screw Cap, completely pierce foil.
- Turn the Safety Screw Connector Clockwise until it is securely fastened.
- Close clamp on set before inverting container.
- Invert container and suspend, using the hanging ring on the bottom of the container.
- **Hang product up to 48 hours when clean technique and only one Screw Cap feeding set is used.**

Instructions for Use: 8-fl-oz-can

- Store unopened at room temperature; avoid extreme temperatures.
- Shake well. Jevity 1.5 Cal is ready to use
- Clean top of container thoroughly before opening.
- Once opened, cover, refrigerate and **use within 48 hours.**

Oral Feeding:

- May be used for total or supplemental nutrition.
- May be fed at room temperature or chilled.

Tube Feeding:

- Follow physician's instructions.
- Adjust flow rate and volume according to patient's condition and tolerance.
- Feed at room temperature.
- Pump feeding is recommended; use an 10 Fr or larger tube. For gravity feeding use a 12 Fr or larger tube.
- Additional fluid requirements should be met by giving water between or after feedings or when flushing the tube.
- Avoid contamination during preparation and use.