

Product Information: Similac® PM 60/40

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Similac® PM 60/40

Low-Iron Infant Formula

- For infants who would benefit from lowered mineral intake, including those with impaired renal function.
- Use under medical supervision.
- Mineral levels closely approximating the mineral content of human milk (60:40 ratio of whey to casein).
- Calcium-to-phosphorus ratio and content designed to treat serum calcium disorders—both hypercalcemia and hypocalcemia due to hyperphosphatemia.
- Gluten-free.
- Halal.
- Kosher.

Safety Precautions

- In conditions where the infant is losing abnormal quantities of one or more electrolytes, it may be necessary to supply electrolytes from sources other than the formula.
- Additional iron should be supplied from other sources.
- Never use a microwave to warm formula. Serious burns can result.
- Not for IV use.



Ingredients

Unflavored Powder:

Lactose, High Oleic Safflower Oil, Whey Protein Concentrate, Soy Oil, Coconut Oil, Sodium Caseinate. Less than 2% of the Following: Potassium Citrate, Calcium Phosphate, Calcium Carbonate, Magnesium Chloride, Potassium Chloride, Ascorbic Acid, m-Inositol, Salt, Choline Chloride, Taurine, Ascorbyl Palmitate, Ferrous Sulfate, Zinc Sulfate, Mixed Tocopherols, L-Carnitine, Niacinamide, d-Alpha-Tocopheryl Acetate, Calcium Pantothenate, Cupric Sulfate, Vitamin A Palmitate, Thiamine Chloride Hydrochloride, Riboflavin, Pyridoxine Hydrochloride, Beta-Carotene, Folic Acid, Manganese Sulfate, Potassium Iodide, Phylloquinone, Biotin, Sodium Selenate, Vitamin D3, Cyanocobalamin, and Potassium Hydroxide.

Allergens: Contains milk ingredients.

Availability

List Number	Item
00850	Similac PM 60/40 Powder / 14.1-oz (400-g) can / 6 ct

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Nutrition Information - Unflavored Powder

	100 Cal	1000 mL
	Value	Value
Osmolality, mOsm/kg H ₂ O	280	280
Calories	100	676
Volume, mL	148	1000
Protein, g	2.2	15
Fat, g	5.6	37.9
Carbohydrate, g	10.2	69.0
Water, g	134	899
Linoleic Acid, mg	1000	6763
Potential Renal Solute Load, mOsm	18.3	124.1
Vitamin A, IU	300	2029
Vitamin D, IU	60	406
Vitamin E, IU	1.5	10.1
Vitamin K, mcg	8	54
Thiamin (Vitamin B1), mcg	100	676
Riboflavin (Vitamin B2), mcg	150	1014
Vitamin B6, mcg	60	406
Vitamin B12, mcg	0.3	1.7
Niacin, mcg	1050	7101
Folic Acid (Folacin), mcg	15	101
Pantothenic Acid, mcg	450	3043
Biotin, mcg	5	30.4
Vitamin C (Ascorbic Acid), mg	9	61
Choline, mg	12	81
Inositol, mg	24	162
Calcium, mg	56	379
Calcium, mEq	2.8	18.9
Phosphorus, mg	28	189
Magnesium, mg	6	40.6
Iron, mg	0.7	4.7
Zinc, mg	0.8	5.1
Manganese, mcg	5	34
Copper, mcg	90	609
Iodine, mcg	6	41
Selenium, mcg	1.8	12.2
Sodium, mg	24	162
Potassium, mg	80	541
Chloride, mg	59	399

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Preparation

Directions for Preparation and Use

Your baby's health depends on carefully following these easy directions. Proper hygiene, handling and storage are important when preparing infant formula. Failure to follow these directions could result in severe harm. Ask your baby's doctor if you need to boil (sterilize) bottles, nipples and rings before use.

- Wash your hands, surfaces and utensils.
- Pour desired amount of water into a clean bottle (see mixing guide).
- Add 1 unpacked level scoop (8.7 g) to each 2 fl oz of water
- Return dry scoop to can.
- Cap bottle; shake well; attach nipple.
- Once feeding begins, **use within 1 hour or discard.**

Powder Mixing Guide

Measure water	+	Add scoop(s) of <i>unpacked level powder</i> using enclosed scoop	=	Finished bottle (approx)
2 fl oz		1 scoop (8.7 g)		2 fl oz
4 fl oz		2 scoops		4 fl oz
6 fl oz		3 scoops		6 fl oz

For a larger amount of formula, add 1 unpacked, level cup of powder (100 g) to 23 fl oz of water (makes approximately 26 fl oz of formula).

When mixed as directed, the contents of this can will make approximately 102 fl oz of formula.

Storage & Handling

Powder

- Once mixed, store bottles in refrigerator and **feed to baby within 24 hours.**
- Store unopened or opened container at room temperature; avoid extreme temperatures.
- **Use opened can contents within 1 month.**