


Product Information: Calcilo XD®

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Calcilo XD®

Low-Calcium/Vitamin D-Free Infant Formula With Iron

- 1 Nutrition support of infants with hypercalcemia, as may occur in infants with Williams syndrome, osteopetrosis, and primary neonatal hyperparathyroidism, and when a low-calcium, vitamin D-free formula is needed.
- 1 Use under medical supervision.
- 1 No vitamin D.
- 1 The only commercially available formula for the management of hypercalcemia in infants.
- 1 Nutritionally complete—when prepared as directed, meets the American Academy of Pediatrics Committee on Nutrition recommendations for all nutrients including vitamins and minerals except calcium, vitamin D and phosphorus.
- 1 Only a minimal amount of calcium (15 mg/100 g powder; 2.9 mg/100 Cal).
- 1 L-carnitine (7 mg/100 g) and taurine (35 mg/100 g).
- 1 Eliminates the need to prepare a modular formula.
- 1 Kosher.
- 1 Halal.
- 1 Lactose-free.
- 1 Gluten-free.



Safety Precautions

- 1 Must be given only to infants whose calcium and/or vitamin D intake must be severely restricted, as in hypercalcemia, and who are under strict medical supervision. Calcium, vitamin D and phosphorus requirements should be determined by appropriate laboratory tests. This formula does not supply the minimum recommended intake of calcium, vitamin D or phosphorus for normal infants.
- 1 Not for IV use.
- 1 **Never use a microwave oven to warm formula.** Serious burns can result.
- 1 Powdered infant formulas are not sterile and should not be fed to premature infants or infants who might have immune problems unless directed and supervised by your baby's doctor.

Ingredients

Powder Unflavored:

Corn Syrup, Coconut Oil, Corn Oil, Whey Protein Concentrate, Sodium Caseinate. Less than 1.0% of: Potassium Phosphate, Magnesium Chloride, Potassium Chloride, Potassium Citrate, Ascorbic Acid, m-Inositol, Salt, Choline Chloride, Ferrous Sulfate, Potassium Hydroxide, Taurine, Zinc Sulfate, L-Carnitine, d-Alpha-Tocopheryl Acetate, Niacinamide, Calcium Pantothenate, Vitamin A Palmitate, Cupric Sulfate, Thiamine Chloride Hydrochloride, Riboflavin, Pyridoxine Hydrochloride, Folic Acid, Manganese Sulfate, Potassium Iodide, Phylloquinone, Biotin, Sodium Selenate, and Cyanocobalamin.

Allergens: Contains milk ingredients.

Availability

List Number	Item
53328	Calcilo XD / 13.2 oz (375 g) Can / 6 ct

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Abbott Nutrition Abbott Laboratories
Columbus, OH 43219-3034
1-800-227-5767

2 of 4



Nutrition Information - Powder Unflavored

	100 Cal (5 fl oz, prepared as directed)		100 g Powder	
	Value	%RDI / %DV	Value	%RDI / %DV
Volume, mL	148			
Protein, g	2.2		11.4	
Fat, g	5.6		28.7	
Carbohydrate, g	10.2		52.3	
Water, g	134		2.5	
Linoleic Acid, mg	1300		6660	
Calories	100		513	
Potential Renal Solute Load, mOsm	18		94	
Vitamin A, IU	300		1540	
Vitamin D, IU	0		0	
Vitamin E, IU	1.9		10	
Vitamin K, mcg	8		41	
Thiamin (Vitamin B1), mcg	100		513	
Riboflavin (Vitamin B2), mcg	150		770	
Vitamin B6, mcg	60		310	
Vitamin B12, mcg	0.3		1.3	
Niacin, mcg	1050		5400	
Folic Acid (Folacin), mcg	15		77	
Pantothenic Acid, mcg	450		2300	
Biotin, mcg	4.5		23	
Vitamin C, mg	9		46	
Choline, mg	12		62	
Inositol, mg	24		123	
Calcium, mg	2.9		15	
Phosphorus, mg	25		128	
Magnesium, mg	6		31	
Iron, mg	1.8		9.2	
Zinc, mg	0.8		3.8	
Manganese, mcg	5		26	
Copper, mcg	90		460	
Iodine, mcg	6		31	
Selenium, mcg	2		10	
Sodium, mg	24		125	
Sodium, mEq	1.0		5.4	
Potassium, mg	82		420	
Potassium, mEq	2.1		10.7	
Chloride, mg	57		292	
Chloride, mEq	1.6		8.2	

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Abbott Nutrition Abbott Laboratories
 Columbus, OH 43219-3034
 1-800-227-5767



Preparation

Directions for Preparation and Use: Use Only As Directed by a Physician

Your baby's health depends on carefully following these directions. Proper hygiene, handling and storage are important when preparing infant formula. Failure to follow these directions could result in severe harm. Ask your baby's doctor if you need to use cooled, boiled water for mixing and if you need to boil (sterilize) bottles, nipples and rings before use.

- 1 Wash your hands, surfaces and utensils.
- 1 Pour prescribed amount of water into a clean bottle (see mixing guide).
- 1 Add 1 unpacked level scoop (8.6 g) to each 2 fl oz of water
- 1 Return dry scoop to can.
- 1 Cap bottle; shake well; attach nipple.
- 1 Once feeding begins, **use within 1 hour or discard.**

Mixing Guide

Measure Water	Add scoop(s) of <i>unpacked level powder</i> using enclosed scoop	Finished Bottle (approx)
2-fl-oz	1 scoop (8.6 g)	2 fl oz
4-fl-oz	2 scoops	4 fl oz
6-fl-oz	3 scoops	6 fl oz

- 1 To make a larger amount of formula at standard measuring cup, add **1 unpacked, level cup (NOT the enclosed scoop) of powder (105 g)** to 24 fl oz of water. Yields approximately 26 fl oz of formula.
- 1 To maintain freshness, pour prepared formula into individual feeding bottles, cap and store in refrigerator.
- 1 Prepared formula should not be left unrefrigerated.
- 1 When mixed as directed, the contents of one can (375 g) will make approximately 96 fl oz of formula at 20 Cal/fl oz.

Storage & Handling

Powder

- 1 Once mixed, store bottles or container in refrigerator and **feed to baby or child within 24 hours**
- 1 Store unopened or opened can at room temperature; avoid extreme temperatures.
- 1 **Use opened can contents within 1 month.**