

## Product Information: Jevity® 1.2 Cal

For more information, contact your Abbott Nutrition Representative or visit  
[www.abbottnutrition.com](http://www.abbottnutrition.com)

# Jevity® 1.2 Cal

## Complete, Balanced Nutrition® with Fiber

- 1 JEVITY 1.2 CAL is high-protein, fiber-fortified therapeutic nutrition that provides complete, balanced nutrition for long- or short-term tube feeding.
- 1 For tube feeding.
- 1 For supplemental or sole-source nutrition.
- 1 May be used for oral feeding of patients with altered taste perception.
- 1 Use under medical supervision.
- 1 Concentrated calories (1.2 Cal/mL) and high in protein (18.5% of Cal) to help tube-fed patients gain and maintain healthy weight. Protein is needed for lean body tissue maintenance and repair.
- 1 17 g total dietary fiber/L (9 g of a unique fiber blend and 8 g of NutraFlora® scFOS®\*) to help meet daily fiber needs. Fiber helps maintain bowel function.
- 1 Includes 1.9 g Nutraflora scFOS\* /8 fl oz (8 g/1000 mL and 12 g/1500 mL). scFOS are prebiotics that stimulate the growth of beneficial bacteria in the colon.
- 1 Meets or exceeds 100% of the RDIs for 25 essential vitamins and minerals in 1500 Cal (1250 mL).
- 1 Kosher.
- 1 Halal.
- 1 Gluten-free.
- 1 Suitable for lactose intolerance.



## Safety Precautions

- 1 Not for IV use.
- 1 Not for people with galactosemia.

\* NutraFlora® and scFOS® are not registered trademarks of Abbott Laboratories.

## Ingredients

### Liquid Unflavored:

Water, Corn Maltodextrin, Corn Syrup Solids, Sodium Caseinate, Soy Protein Isolate, Canola Oil, Corn Oil, Short-chain Fructooligosaccharides, Medium chain Triglycerides, Calcium Caseinate, Soy Fiber. Less than 0.5% of the Following: Oat Fiber, Potassium Citrate, Calcium Phosphate, Magnesium Phosphate, Gum Arabic, Potassium Chloride, Soy Lecithin, Cellulose Gum, Choline Chloride, Ascorbic Acid, Salt, Potassium Hydroxide, L-Carnitine, Taurine, dl-Alpha-Tocopheryl Acetate, Ferrous Sulfate, Sodium Citrate, Zinc Sulfate, Manganese Sulfate, Niacinamide, Copper Sulfate, Calcium Pantothenate, Thiamine Hydrochloride, Riboflavin, Pyridoxine Hydrochloride, Beta-Carotene, Vitamin A Palmitate, Chromium Chloride, Folic Acid, Sodium Molybdate, Potassium Iodide, Sodium Selenate, Biotin, Phylloquinone, Vitamin D3, and Vitamin B12.

**Allergens:** Contains milk and soy ingredients.

## Availability

List Number	Item
62667	Jevity 1.2 Cal / 50.7 fl oz (1.5 L) Ready-To-Hang Prefilled Container / 6 ct
62683	Jevity 1.2 Cal / 33.8 fl oz (1 L) Ready-To-Hang Prefilled Container / 8 ct
64625	Jevity 1.2 Cal / 8 fl oz (237 mL) Recloseable Carton / 24 ct

For more information, contact your Abbott Nutrition Representative or visit  
[www.abbottnutrition.com](http://www.abbottnutrition.com)

Abbott Nutrition Abbott Laboratories  
Columbus, OH 43219-3034  
1-800-227-5767

2 of 5



### Nutrition Information - Liquid Unflavored

	8 fl oz (237 mL)		1 L (33.8 fl oz)		1.5 L (50.7 fl oz)	
	Value	%RDI / %DV	Value	%RDI / %DV	Value	%RDI / %DV
Protein, g	13.2		55.5		83.3	
Fat, g	9.3		39.3		59	
Carbohydrate, g	40.2		169.4		254	
Dietary Fiber, g	4*		17*		25.5*	
Total Sugars, g	3		13		20	
Added Sugars, g	3		13		20	
L-Carnitine, mg	36		150		225	
Taurine, mg	36		150		225	
Water, g(mL)	191		807		1211	
Calories	285		1200		1800	
Vitamin A, mcg RAE	171		720		1080	
Beta-Carotene, mcg	307		1295		1943	
Vitamin D, mcg	3.8		16		24	
Vitamin E, mg	9.2		38		57	
Vitamin K, mcg	29		123		185	
Vitamin C, mg	59		250		375	
Folate, mcg DFE	76		320		480	
Folic Acid, mcg	46		192		288	
Thiamin (Vitamin B1), mg	0.41		1.7		2.6	
Riboflavin (Vitamin B2), mg	0.59		2.5		3.7	
Vitamin B6, mg	0.33		1.4		2.1	
Vitamin B12, mcg	2.3		9.6		14.4	
Niacin, mg NE	4		17		26	
Choline, mg	142		600		900	
Biotin, mcg	5.7		24		36	
Pantothenic Acid, mg	0.95		4		6.0	
Sodium, mg	253		1067		1600	
Sodium, mEq	11		46.4		69.6	
Potassium, mg	566		2390		3585	
Potassium, mEq	14.5		67.4		91.7	
Chloride, mg	356		1500		2250	
Chloride, mEq	10		42.4		63.6	
Calcium, mg	284		1200		1800	
Phosphorus, mg	284		1200		1800	
Magnesium, mg	88		370		555	
Iodine, mcg	36		150		225	
Manganese, mg	0.81		3.4		5.1	
Copper, mg	0.41		1.7		2.6	
Zinc, mg	2.5		11		16	

For more information, contact your Abbott Nutrition Representative or visit  
[www.abbottnutrition.com](http://www.abbottnutrition.com)

### Nutrition Information - Liquid Unflavored

	8 fl oz (237 mL)		1 L (33.8 fl oz)		1.5 L (50.7 fl oz)	
	Value	%RDI / %DV	Value	%RDI / %DV	Value	%RDI / %DV
Iron, mg	3.4		14.4		22	
Selenium, mcg	14		58		87	
Chromium, mcg	14		58		87	
Molybdenum, mcg	23		99		148	

### Liquid Unflavored Footnotes & References

#### **Per 8 fl oz (237 mL)**

\*2.1 g of total dietary fiber from a unique fiber blend and 1.9 g of short-chain fructooligosaccharides.

†795 IU of vitamin A activity supplied by 0.60 mg beta-carotene.

#### **Per 1 L (33.8 fl oz)**

\*9 g of total dietary fiber from a unique fiber blend and 8 g of short-chain fructooligosaccharides.

#### **Per 1.5 L (50.7 fl oz)**

\*13.5 g of total dietary fiber from a unique fiber blend and 12 g of short-chain fructooligosaccharides.

### Preparation

#### Instructions for Use:

##### 8 fl oz carton

- 1 Store unopened at room temperature; avoid extreme temperatures.
- 1 Shake well prior to opening.
- 1 Once opened, reclose, refrigerate and **use within 48 hours**.

#### Oral Feeding:

- 1 May be used for total or supplemental nutrition.
- 1 May be fed at room temperature or chilled.

#### Tube Feeding:

- 1 Follow physician's instructions.
- 1 Adjust flow rate and volume according to patient's condition and tolerance.
- 1 Feed at room temperature using a feeding pump or syringe.
- 1 Additional fluid requirements should be met by giving water between or after feedings or when flushing the tube.
- 1 Avoid contamination during preparation and use.

#### **Ready-To-Hang Container**

All liquid medical foods, regardless of type of administration system, require careful handling because they can support microbial growth. Follow these instructions for clean technique and proper setup to reduce the potential for microbial contamination.

NOTE: Failure to follow the Instructions for Use increases the potential for microbial contamination and reduces hangtime.

- 1 Administer product at room temperature.
- 1 THOROUGHLY wash hands with soap and water before handling container or feeding set.
- 1 Turn container upside down and SHAKE VIGOROUSLY, using a twisting motion for at least 10 seconds.
- 1 DO NOT touch any part of the container or feeding set that comes into contact with the formula.
- 1 When initiating feeding, follow physician's instructions. Adjust flow rate and volume according to patient's condition and tolerance.
- 1 Additional fluid requirements should be met by giving water between or after feedings or when flushing the tube.

#### For Use with Enteral Feeding Pumps:

- 1 Remove dust cover from RTH Safety Screw Cap.
- 1 Remove dust cover from feeding set connector.
- 1 Insert feeding set connector into port of RTH Safety Screw Cap and completely pierce foil.
- 1 Turn connector collar clockwise until it is securely fastened.
- 1 Close clamp on set before inverting container.
- 1 Invert container and suspend, using hanging ring on bottom of container.
- 1 Follow directions for use provided by manufacturer of feeding sets.
- 1 Unless a shorter hang time is specified by the set manufacturer, hang product for up to 48 hours after initial connection when clean technique and only one new set are used. Otherwise hang for no more than 24 hours.