

Product Information: LiquiProtein®

For more information, contact your Abbott Nutrition sales representative or visit
www.abbottnutrition.ca

Usage

- 1 For use as a modular protein supplement to be administered through a patient's feeding tube.
- 1 Suggested indications:
 - Protein-energy malnutrition;
 - Cancer;
 - Wounds/pressure injuries;
 - Pre- and post-surgery;
 - Trauma/burns.
- 1 Use under medical supervision.

Features

- 1 A commercially sterile, extensively hydrolyzed liquid protein modular. Eliminates the need for protein powder mixing to reduce the risk of contamination.¹
- 1 Allows for customized protein intake.
- 1 Lactose-free.
- 1 Low-residue.
- 1 Gluten-free.

Safety Precautions

- 1 Not for use as a sole source of nutrition.
- 1 If signs of intolerance develop, slow administration rate or discontinue.
- 1 Not for parenteral use.
- 1 Not suitable for people with galactosemia.

¹ Bankhead R, et al. *JPEN*. 2009;33(2):122-67.

Ingredients

Liquid Unflavored:

Water, and Casein Hydrolysate (Milk).

Allergens: Contains: Milk

Availability

List Number	Item
6231713	LiquiProtein®; 54-mL plastic bottles; 48/case



Nutrition Information - Liquid Unflavored

	6 mL	54 mL
	Value	Value
Energy, Cal (kJ)	4 (17)	35 (149)
Protein, g	1	9
Fat, g	0	0
Carbohydrate, g	0	0
Sodium, mg	5	75
Potassium, mg	10	100
Iron, mg	0	0.1

Product Characteristics

	Value
Caloric Density (Cal/mL)	0.67
Protein (% of total energy)	100
Protein Source	Casein hydrolysate
Fat (% of total energy)	0
MCT oil (% of total fats)	N/A
Omega-6:Omega-3 Ratio	N/A
Fat Source	N/A
Carbohydrate (% of total energy)	0
Carbohydrate Source	N/A
Total Cal:g Nitrogen Ratio	25:1
Nonprotein Cal:g Nitrogen Ratio	N/A
Water (g/L)	810
Osmolality (mOsm/kg water)	750 (when mixed 1:1 with water)
Renal Solute Load (mOsm/L)	N/A
Recommended Tube Size for Gravity/Pump Feeding (Fr)	N/A

Liquid Unflavored Footnotes & References

Per 6 mL

*5% or less is a little, 15% or more is a lot. Not a significant source of saturated fat, trans fat, fibre, sugars, cholesterol, calcium and iron. Other inherent minerals may be present in trace amounts.

Per 54 mL

*5% or less is a little, 15% or more is a lot. Not a significant source of saturated fat, trans fat, fibre, sugars, cholesterol, calcium and iron. Other inherent minerals may be present in trace amounts.

Preparation

Directions for Use:

- 1 For use in gastric feeding tube: no dilution is required.
- 1 For use in small bowel feeding tube: dilute with an equal part of water.

1. Administer by syringe through the feeding tube. Flush with 20-30 mL of water[†] before and after administration.

[†] Adjust flush volume depending on fluid requirements.

Storage & Handling

Directions for Use:

- 1 Store unopened at room temperature; avoid extreme temperatures.
- 1 Once opened, refrigerate and use within 72 hours.
- 1 Cover prepared, unused mixture immediately and use within 24 hours.