

Product Information: PediaSure® 1.5 Cal with Fiber

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

PediaSure® 1.5 Cal with Fiber

Complete, Balanced Nutrition®



- 1 PEDIASURE 1.5 CAL WITH FIBER is a higher* caloric density product designed to meet the higher energy requirements of pediatric patients who are at risk for malnutrition, require a higher caloric density, or have fluid restrictions. It provides a source of complete, balanced nutrition for children 1 to 13 years of age.
- 1 For oral or tube feeding.
- 1 May be used as a sole source of nutrition or as a supplement.
- 1 Use under medical supervision.
- 1 Meets or exceeds 100% of the DRIs for protein and 25 essential vitamins and minerals
 - in 1000 mL for children 1 to 8 years of age.
 - in 1500 mL for children 9 to 13 years of age.
- 1 Excellent source of high-quality protein designed to meet the acceptable macronutrient distribution range of protein for children 1 to 13 years of age.
- 1 Vitamin D, calcium and phosphorus for strong bones.
- 1 Prebiotics designed to support digestive tract health^{1,2,3,4} and antioxidants (vitamins C & E and selenium) to support the immune system.
- 1 DHA (32 mg per 8 fl oz) and lutein to support the brain and eyes.^{5,6}
- 1 No artificial growth hormones.[†]
- 1 Kosher.
- 1 Halal.
- 1 Gluten-free.
- 1 Suitable for lactose intolerance.

Safety Precautions

- 1 Not intended for infants under 1 year of age unless specified by a physician.
- 1 Not for IV use.
- 1 Not for children with galactosemia.

* Caloric value is 1.5 Cal/mL compared to PediaSure Enteral Formula 1.0 Cal at 1.0 Cal/mL.

† No significant difference has been shown between milk derived from rbST-treated and non-rbST-treated cows.

¹ Bornet FR, et al. *Nutr Rev* 2002;60:326-334.

² Tokunaga T, et al. *Bifidus* 1993;6:143-150.

³ Hidaka H, et al. *Bifidobact Microflora* 1986;5:37-50.

⁴ Bouhnik Y, et al. *Nutr J* 2006; doi:10.1186/1475-2891-5-8.

⁵ Lauritzen L, et al. *Prog Lipid Res.* 2001;40:1-94.

⁶ Krinsky NI, et al. *Annu Rev Nutr.* 2003;23:171-201.

Ingredients

Liquid Vanilla:

Water, Corn Maltodextrin, Blend of Vegetable Oils (High Oleic Safflower, Soy, Medium Chain Triglycerides), Milk Protein Concentrate, Soy Protein Isolate, Short-chain Fructooligosaccharides. Less than 0.5% of: Vitamins & Minerals (Potassium Citrate, Potassium Chloride, Magnesium Phosphate, Calcium Phosphate, Ascorbic Acid, Choline Chloride, Calcium Carbonate, Potassium Phosphate, Ferrous Sulfate, dl-Alpha-Tocopheryl Acetate, Zinc Sulfate, Niacinamide, Calcium Pantothenate, Manganese Sulfate, Thiamine Hydrochloride, Pyridoxine Hydrochloride, Riboflavin, Vitamin A Palmitate, Copper Sulfate, Folic Acid, Chromium Chloride, Potassium Iodide, Sodium Molybdate, Sodium Selenate, Biotin, Phylloquinone, Vitamin D3, Vitamin B12), Oat Fiber, Natural & Artificial Flavor, Soy Fiber, Cellulose Gel, Soy Lecithin, Tuna Oil, Monoglycerides, Cellulose Gum, Potassium Hydroxide, Carrageenan, Inositol, Taurine, Sucralose, Salt, L-Carnitine, and Lutein.

Allergens: Contains milk and soy ingredients.

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Abbott Nutrition Abbott Laboratories
Columbus, OH 43219-3034
1-800-227-5767

2 of 5



PediaSure® 1.5 Cal with Fiber

Complete, Balanced Nutrition®

Availability

List Number	Item
56411	PediaSure 1.5 Cal w Fiber Vanilla / 8 fl oz (237 mL) Can / 24 ct (DISC)
62749	PediaSure 1.5 Cal w Fiber Vanilla / 33.8 fl oz (1 L) RTH Prefilled Container / 8 ct (DISC)
67374	PediaSure 1.5 Cal w Fiber Vanilla / 8 fl oz (237 mL) Can / 24 ct
67376	PediaSure 1.5 Cal w Fiber Vanilla / 33.8 fl oz (1 L) Ready-To-Hang Prefilled Container / 8 ct

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Abbott Nutrition Abbott Laboratories
Columbus, OH 43219-3034
1-800-227-5767



PediaSure® 1.5 Cal with Fiber

Complete, Balanced Nutrition®

Nutrition Information - Liquid Vanilla

	8 fl oz (237 mL)		1000 mL (1.1 QT)	
	Value	%RDI / %DV	Value	%RDI / %DV
Protein, g	14		59	
Fat, g	16		68	
Saturated Fat, g	3.5		14	
Trans Fat, g	0		0	
Polyunsaturated Fat, g	5		20	
Monounsaturated Fat, g	7		30	
Cholesterol, mg	10		40	
Carbohydrate, g	39		166	
Dietary Fiber, g	3		13	
scFOS®, g	1.7		7	
Total Sugars, g	3		11	
Added Sugars, g	3		10	
L-Carnitine, mg	4.0		17	
Taurine, mg	18		76	
Water, g	185		780	
Calories	350		1500	
Vitamin A, mcg RAE	140		590	
Vitamin D, mcg	6.0		25	
Vitamin E, mg	3.0		13	
Vitamin K, mcg	18		76	
Vitamin C, mg	23		97	
Folate, mcg DFE	60		253	
Folic Acid, mcg	36		152	
Thiamin (Vitamin B1), mg	0.30		1.3	
Riboflavin (Vitamin B2), mg	0.33		1.4	
Vitamin B6, mg	0.34		1.4	
Vitamin B12, mcg	0.48		2.0	
Niacin, mg NE	3.2		14	
Choline, mg	80		338	
Biotin, mcg	8.0		34	
Pantothenic Acid, mg	1.3		5.5	
Inositol, mg	20		84	
Sodium, mg	90		380	
Potassium, mg	470		1980	
Chloride, mg	230		970	
Calcium, mg	330		1390	
Phosphorus, mg	250		1050	
Magnesium, mg	40		169	
Iodine, mcg	23		97	

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Nutrition Information - Liquid Vanilla

	8 fl oz (237 mL)		1000 mL (1.1 QT)	
	Value	%RDI / %DV	Value	%RDI / %DV
Manganese, mg	0.46		1.9	
Copper, mg	0.14		0.59	
Zinc, mg	1.7		7.2	
Iron, mg	2.7		11	
Selenium, mcg	8.0		34	
Chromium, mcg	9.0		38	
Molybdenum, mcg	9.0		38	

Preparation

Instructions for Use:

Can

- 1 Shake well.
- 1 Store unopened at room temperature.
- 1 Clean top of container thoroughly before opening.
- 1 After opening, refrigerate and **use within 48 hours**.
- 1 Tube Feeding
 - 1 Follow physician's instructions for flow rate, volume, dilution and need for additional fluids.
- 1 Oral Feeding
 - 1 Best Served Chilled

Ready-To-Hang Container

All liquid medical foods, regardless of type of administration system, require careful handling because they can support microbial growth. Follow these instructions for clean technique and proper setup to reduce the potential for microbial contamination.

NOTE: Failure to follow the **Instructions for Use** increases the potential for microbial contamination and reduces hangtime.

- 1 Administer product at room temperature.
- 1 THOROUGHLY wash hands with soap and water before handling container or feeding set.
- 1 Turn container upside down and SHAKE VIGOROUSLY, using a twisting motion for at least 10 seconds.
- 1 DO NOT touch any part of the container or feeding set that comes into contact with the formula.
- 1 When initiating feeding, follow physician's instructions. Adjust flow rate and volume according to patient's condition and tolerance.
- 1 Additional fluid requirements should be met by giving water between or after feedings or when flushing the tube.
- 1 Always follow directions for use provided with feeding set.
- 1 For Use with Enteral Feeding Set:
 - 1 Remove dust cover from RTH Safety Screw Cap.
 - 1 Remove dust cover from feeding set connector.
 - 1 Insert feeding set connector into port of RTH Safety Screw Cap and completely pierce foil.
 - 1 Turn connector collar clockwise until it is securely fastened.
 - 1 Close clamp on feeding set before inverting container.
 - 1 Invert container and suspend, using hanging ring on the bottom of container.
 - 1 Unless a shorter hang time is specified by the set manufacturer, hang product for up to 48 hours after initial connection when clean technique and only one new set are used. Otherwise hang for no more than 24 hours.

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com