

Product Information: PediaSure® Enteral Formula 1.0 Cal with Fiber

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

PediaSure® Enteral Formula 1.0 Cal with Fiber

Complete, Balanced Nutrition®

- 1 PEDIASURE ENTERAL FORMULA 1.0 CAL WITH FIBER is a source of Complete, Balanced Nutrition® especially designed for tube feeding children 1 to 13 years of age.
- 1 May be used as the sole source of nutrition or as a supplement.
- 1 Use under medical supervision.
- 1 1.0 Cal/mL, 240 Cal/8 fl oz, from a balanced distribution of protein, fat, and carbohydrate.
- 1 Provides 100% or more of the Dietary Reference Intakes (DRIs) for protein and 25 essential vitamins and minerals in 1000 mL for children 1 to 8 years of age, and in 1500 mL for children 9 to 13 years of age.
- 1 Vitamin D, calcium and phosphorus for strong bones.
- 1 DHA omega-3* to help support brain and eye development.^{1,2}
- 1 Prebiotics designed to support digestive tract health^{3,4,5} and antioxidants (vitamins C & E and selenium) to support the immune system.
- 1 No artificial growth hormones.[†]
- 1 Kosher.
- 1 Halal.
- 1 Gluten-free.
- 1 Suitable for lactose intolerance.



Safety Precautions

- 1 Not intended for infants under 1 year of age unless specified by a physician.
- 1 Not for IV use.
- 1 Not for children with galactosemia.

* 32 mg DHA per 8 fl oz serving.

† Milk ingredients are derived from milk of non-rbST-treated cows, which is not different from milk of rbST-treated cows.

¹ Lauritzen L, et al. *Prog Lipid Res.* 2001;40:1-94.

² Krinsky NI, et al. *Annu Rev Nutr.* 2003;23:171-201.

³ Bornet FR, et al. *Nutr Rev.* 2002;60(11):326-334.

⁴ Tokunaga T, et al. *Bifidus.* 1993;6(2):143-150.

⁵ Hidaka H, et al. *Bifidobacteria Microflora.* 1986;5:37-50.

Ingredients

Liquid Vanilla:

Water, Corn Maltodextrin, Blend of Vegetable Oils (High Oleic Safflower, Soy, Medium Chain Triglycerides), Milk Protein Concentrate, Sugar, Soy Protein Isolate, Short-chain Fructooligosaccharides. Less than 0.5% of: Vitamin & Minerals (Potassium Phosphate, Calcium Carbonate, Potassium Chloride, Potassium Citrate, Calcium Phosphate, Ascorbic Acid, Choline Chloride, Ferrous Sulfate, dl-Alpha-Tocopheryl Acetate, Zinc Sulfate, Niacinamide, Calcium Pantothenate, Manganese Sulfate, Thiamine Hydrochloride, Pyridoxine Hydrochloride, Riboflavin, Vitamin A Palmitate, Copper Sulfate, Magnesium Phosphate, Folic Acid, Chromium Chloride, Potassium Iodide, Sodium Molybdate, Sodium Selenate, Biotin, Phylloquinone, Vitamin D3, Vitamin B12), Oat Fiber, Soy Fiber, Natural & Artificial Flavors, Gum Arabic, Cellulose Gel, Tuna Oil*, Cellulose Gum, Soy Lecithin, Monoglycerides, Carrageenan, Salt, Potassium Hydroxide, Inositol, Taurine, L-Carnitine, and Lutein.

* Source of docosahexaenoic acid (DHA).

Allergens: Contains milk and soy ingredients.

Availability

List Number	Item
67403	PediaSure Enteral 1.0 Cal w Fiber Vanilla / 8 fl oz (237 mL) Can / 24 ct
67405	PediaSure Enteral 1.0 Cal w Fiber Vanilla / 33.8 fl oz (1 L) Ready-To-Hang Prefilled Container / 8 ct

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Abbott Nutrition Abbott Laboratories
Columbus, OH 43219-3034
1-800-227-5767



PediaSure® Enteral Formula 1.0 Cal with Fiber

Complete, Balanced Nutrition®

Nutrition Information - Liquid Vanilla

	8 fl oz (237 mL)		33.8 fl oz (1 L)	
	Value	%RDI / %DV	Value	%RDI / %DV
Volume, mL	237			
Protein, g	7		30	
Fat, g	9			
Saturated Fat, g	2.5		11	
Trans Fat, g	0		0	
Polyunsaturated Fat, g	2.5		11	
Monounsaturated Fat, g	3.5		15	
Cholesterol, mg	5		21	
Carbohydrate, g	34			
Dietary Fiber, g	3		13	
scFOS, g	1.7		7	
Total Sugars, g	7		30	
Added Sugars, g	7		30	
L-Carnitine, mg	4.0		16.9	
Taurine, mg	18		76	
Water, g	200		844	
Calories	240		1000	
Vitamin A, mcg RAE	140		590	
Vitamin D, mcg	6.0		25	
Vitamin E, mg	3.0		13	
Vitamin K, mcg	18		76	
Vitamin C, mg	23		97	
Folate, mcg DFE	60		253	
Folic Acid, mcg	36		152	
Thiamin (Vitamin B1), mg	0.30			
Riboflavin (Vitamin B2), mg	0.33			
Vitamin B6, mg	0.34		1.4	
Vitamin B12, mcg	0.48		2.0	
Niacin, mg NE	3.2		14	
Choline, mg	80		338	
Biotin, mcg	8.0		34	
Pantothenic Acid, mg	1.3		5.5	
Inositol, mg	20		84	
Sodium, mg	90		380	
Potassium, mg	470		1983	
Chloride, mg	230		970	
Calcium, mg	330		1392	
Phosphorus, mg	250		1055	
Magnesium, mg	40		169	

For more information, contact your Abbott Nutrition Representative or visit www.abbottnutrition.com

Abbott Nutrition Abbott Laboratories
Columbus, OH 43219-3034
1-800-227-5767



Nutrition Information - Liquid Vanilla

	8 fl oz (237 mL)		33.8 fl oz (1 L)	
	Value	%RDI / %DV	Value	%RDI / %DV
Iodine, mcg	23		97	
Manganese, mg	0.46		1.9	
Copper, mg	0.14		0.59	
Zinc, mg	1.7		7.2	
Iron, mg	2.7		11	
Selenium, mcg	8.0		34	
Chromium, mcg	9.0		38	
Molybdenum, mcg	9.0		38	
Potential Renal Solute Load, mOsm	66			

Preparation

Instructions for Use:

8 fl oz Container

- 1 Shake well.
- 1 Store unopened at room temperature.
- 1 Clean top of container thoroughly before opening.
- 1 After opening, refrigerate and consume product within 48 hours. If consuming product directly from the container, refrigerate and consume within 24 hours.

Tube Feeding

- 1 Follow physician's instructions for flow rate, volume, dilution and need for additional fluids.

Ready-To-Hang Container

All medical foods, regardless of type of administration system, require careful handling because they can support microbial growth. Follow these instructions for clean technique and proper setup to reduce the potential for microbial contamination.

NOTE: Failure to follow the **INSTRUCTIONS FOR USE** increases the potential for microbial contamination and may reduce hangtime.

- 1 Administer product at room temperature.
- 1 THOROUGHLY wash hands with soap and water before handling container or feeding set.
- 1 Turn container upside down and SHAKE VIGOROUSLY, using a twisting motion for at least 10 seconds.
- 1 DO NOT touch any part of the container or feeding set that comes into contact with the formula.
- 1 When initiating feeding, follow physician's instructions. Adjust flow rate and volume according to patient's condition and tolerance.
- 1 Additional fluid requirements should be met by giving water between or after feedings or when flushing the tube.

For Use with Enteral Feeding Pumps:

- 1 Remove dust cover from RTH Safety Screw Cap.
- 1 Remove dust cover from feeding set connector.
- 1 Insert feeding set connector into port of RTH Safety Screw Cap and completely pierce foil.
- 1 Turn connector collar clockwise until it is securely fastened.
- 1 Close clamp on set before inverting container.
- 1 Invert container and suspend, using hanging ring on bottom of container.

Precautions

- 1 Follow directions for use provided by manufacturer of feeding sets.
- 1 Unless a shorter hang time is specified by the set manufacturer, hang product for up to 48 hours after initial connection when clean technique and only one new set are used. Otherwise hang for no more than 24 hours.

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com