

## Product Information: PediaSure® Peptide 1.0 Cal

For more information, contact your Abbott Nutrition Representative or visit  
[www.abbottnutrition.com](http://www.abbottnutrition.com)

# PediaSure® Peptide 1.0 Cal

## Peptide-Based Nutrition for Children

- 1 PEDIASURE PEPTIDE 1.0 Cal is a nutritionally complete, peptide-based formula for the nutritional needs of children ages 1-13 years with malabsorption, maldigestion, and other GI conditions.
- 1 For oral or tube feeding.
- 1 For supplemental or sole-source nutrition.
- 1 Use under medical supervision.
- 1 Nutrient blend designed to help support tolerance and absorption:
  - Hydrolyzed, whey-dominant protein.
  - 60% MCT & structured lipids.
  - Nutraflora® scFOS®\*, a prebiotic fiber.
- 1 Nutrition to help support:
  - Absorption.
  - Tolerance.
  - Growth and development.
- 1 Meets or exceeds 100% of the DRIs for protein and 25 essential vitamins and minerals for children 1 to 8 years of age in 1000 mL (1000 Cal); for children 9 to 13 years of age in 1500 mL (1500 Cal).
- 1 Only peptide-based pediatric product with Kosher status.
- 1 Halal.
- 1 Kosher.
- 1 Gluten-free.
- 1 Suitable for lactose intolerance.



## Safety Precautions

- 1 Not intended for infants under 1 year of age unless specified by a physician.
- 1 Not for IV use.
- 1 Not for children with galactosemia.
- 1 **Feeding sets are for single-patient use only. Use clean technique to avoid set and/or product contamination.**

\* NutraFlora® scFOS® is not a registered trademark of Abbott Laboratories.

## Ingredients

### Liquid Vanilla:

Water, Corn Maltodextrin, Sugar, Whey Protein Hydrolysate, Structured Lipid (Interesterified Canola, Medium-Chain Triglycerides), Medium-Chain Triglycerides, Hydrolyzed Sodium Caseinate. Less than 0.5% of: Short-Chain Fructooligosaccharides, Canola Oil, Calcium Phosphate, Natural & Artificial Flavors, Potassium Citrate, Cellulose Gel, Magnesium Phosphate, Soy Lecithin, Potassium Chloride, Magnesium Chloride, Ascorbic Acid, Carrageenan, Choline Chloride, Sodium Citrate, Cellulose Gum, m-Inositol, Taurine, Sucralose, Ferrous Sulfate, Zinc Sulfate, dl-Alpha-Tocopheryl Acetate, L-Carnitine, Calcium Pantothenate, Niacinamide, Manganese Sulfate, Thiamine Hydrochloride, Pyridoxine Hydrochloride, Riboflavin, Copper Sulfate, Vitamin A Palmitate, Folic Acid, Chromium Chloride, Biotin, Potassium Iodide, Sodium Selenate, Sodium Molybdate, Phylloquinone, Vitamin D3, and Vitamin B12.

**Allergens:** Contains milk and soy ingredients.

## Availability

List Number	Item
62119	PediaSure Peptide 1.0 Ready-to-Drink Vanilla / 8-fl-oz (237-mL) bottle / 24 ct
62121	PediaSure Peptide 1.0 Ready-to-Drink Strawberry / 8-fl-oz (237-mL) bottle / 24 ct
62123	PediaSure Peptide 1.0 Unflavored Ready-to-Drink / 8-fl-oz (237-mL) bottle / 24 ct
62729	PediaSure Peptide 1.0 Unflavored Ready-to-Hang / 1000-mL (1.1-QT) prefilled container / 8 ct

For more information, contact your Abbott Nutrition Representative or visit  
[www.abbottnutrition.com](http://www.abbottnutrition.com)

# PediaSure® Peptide 1.0 Cal

Peptide-Based Nutrition for Children

## Nutrition Information - Liquid Vanilla

	8 fl oz (237 mL)	
	Value	%RDI / %DV
Fat, g	9.6	
Trans Fat, g	0.0	
Carbohydrate, g	31.7	
Dietary Fiber, g	0.7	
scFOS®, g	0.7	
L-Carnitine, mg	4.0	
Taurine, mg	18	
Water, g	201	
Calories	237	
Vitamin A, IU	592	
Vitamin D, IU	237	
Vitamin E, IU	5.4	
Vitamin K, mcg	13	
Vitamin C, mg	24	
Folic Acid, mcg	71	
Thiamin (Vitamin B1), mg	0.6	
Riboflavin (Vitamin B2), mg	0.5	
Vitamin B6, mg	0.6	
Vitamin B12, mcg	1.4	
Niacin, mg	2.4	
Choline, mg	71	
Biotin, mcg	45	
Pantothenic Acid, mg	2.4	
m-Inositol, mg	20	
Sodium, mg	170	
Sodium, mEq	7.4	
Potassium, mg	320	
Potassium, mEq	8.2	
Chloride, mg	240	
Chloride, mEq	6.8	
Calcium, mg	250	
Phosphorus, mg	200	
Magnesium, mg	47	
Iodine, mcg	23	
Manganese, mg	0.4	
Copper, mg	0.2	
Zinc, mg	2.8	
Iron, mg	3.3	
Selenium, mcg	7.6	

For more information, contact your Abbott Nutrition Representative or visit  
[www.abbottnutrition.com](http://www.abbottnutrition.com)

### Nutrition Information - Liquid Vanilla

	8 fl oz (237 mL)	
	Value	%RDI / %DV
Chromium, mcg	7.1	
Molybdenum, mcg	8.5	

### Preparation

#### Instructions for Use: 8-fl-oz Bottle

- 1 **Shake very well.**
- 1 Best served chilled.
- 1 Store unopened at room temperature.
- 1 After opening, refrigerate and **use within 48 hours.**

#### Tube Feeding:

- 1 Follow physician's instructions.
- 1 Adjust flow rate and volume according to child's condition and tolerance.
- 1 Additional fluid may be required.

#### Oral Feeding:

- 1 Best when chilled.

#### Instructions for Use: Ready-To-Hang Container

All medical foods, regardless of type of administration system, require careful handling because they can support microbial growth. Follow these instructions for clean technique and proper setup to reduce the potential for microbial contamination. NOTE: Failure to follow the

**Instructions for Use** increases the potential for microbial contamination and may reduce hangtime.

- 1 Administer product at room temperature.
- 1 THOROUGHLY wash hands with soap and water before handling container or feeding set.
- 1 Turn container upside down and SHAKE VIGOROUSLY, using a twisting motion for at least 10 seconds.
- 1 DO NOT touch any part of the container or feeding set that comes into contact with the formula.
- 1 When initiating feeding, follow physician's instructions. Adjust flow rate and volume according to patient's condition and tolerance.
- 1 Additional fluid requirements should be met by giving water between or after feedings or when flushing the tube.
- 1 Follow directions for use provided by manufacturer of feeding sets.

#### For Use with Enteral Feeding Pumps:

- 1 Remove dust cover from Safety Screw Cap.
- 1 Remove dust cover from feeding set connector.
- 1 Insert feeding set connector into port of RTH Safety Screw Cap and completely pierce foil.
- 1 Turn connector collar clockwise until it is securely fastened.
- 1 Close clamp on feeding set before inverting container.
- 1 Invert container and suspend, using hanging ring on bottom of container.
- 1 Unless a shorter hang time is specified by the set manufacturer, hang product for up to 48 hours after initial connection when clean technique and only one new set are used. Otherwise hang for no more than 24 hours.