

Product Information: PediaSure® Peptide 1.5 Cal

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

PediaSure® Peptide 1.5 Cal

Peptide-Based Nutrition

- 1 PEDIASURE PEPTIDE 1.5 is a nutritionally complete, peptide-based formula for the nutritional needs of children 1 to 13 years of age with malabsorption, maldigestion, and other GI conditions. It is designed to meet the nutritional and caloric needs of kids who require higher caloric density, have fluid restrictions, and are at risk for malnutrition.
- 1 For oral or tube feeding.
- 1 For supplemental or sole-source nutrition.
- 1 Use under medical supervision.
- 1 Complete, Balanced Nutrition®.
- 1 Nutrition to help support absorption, tolerance, and growth and development.
- 1 Nutrient blend designed to help support tolerance and absorption:
 - Prebiotics from Nutraflora® scFOS®.*
 - 60% MCT & structured lipids.
 - Hydrolyzed, whey-dominant protein.
- 1 Meets or exceeds 100% of the DRIs for protein and 25 essential vitamins and minerals for children 1 to 8 years of age in 1000 Cal (667 mL) and for children 9 to 13 years of age in 1500 Cal (1000 mL).
- 1 Antioxidants (vitamins C & E and selenium) to support the immune system.
- 1 Vitamin D (25 mcg/L, 1000 IU/L) and calcium for strong bones.
- 1 Has DHA and lutein to support the brain and eyes.
- 1 Made with milk with no artificial growth hormones.†
- 1 Kosher.
- 1 Halal.
- 1 Gluten-free.
- 1 Suitable for lactose intolerance.



Safety Precautions

- 1 Not intended for infants under 1 year of age unless specified by a physician.
- 1 Not for IV use.
- 1 Not for children with galactosemia.
- 1 **Feeding sets are for single-patient use only. Use clean technique to avoid set and/or product contamination.**

* NutraFlora® and scFOS® are not registered trademarks of Abbott Laboratories.

† No significant difference has been shown between milk derived from rbST-treated and non-rbST-treated cows.

Ingredients

Liquid Vanilla:

Water, Corn Maltodextrin, Structured Lipid (Interesterified Canola Oil & Medium Chain Triglycerides), Whey Protein Hydrolysate, Hydrolyzed Sodium Caseinate, Medium-Chain Triglycerides, Short-chain Fructooligosaccharides. Less than 0.5% of: Vitamins & Minerals (Calcium Phosphate, Potassium Citrate, Potassium Chloride, Potassium Phosphate, Magnesium Phosphate, Calcium Carbonate, Sodium Citrate, Ascorbic Acid, Choline Chloride, Ferrous Sulfate, Magnesium Chloride, Zinc Sulfate, dl-Alpha-Tocopheryl Acetate, Niacinamide, Calcium Pantothenate, Manganese Sulfate, Thiamine Hydrochloride, Pyridoxine Hydrochloride, Riboflavin, Copper Sulfate, Vitamin A Palmitate, Folic Acid, Biotin, Potassium Iodide, Chromium Chloride, Sodium Selenate, Sodium Molybdate, Phylloquinone, Vitamin D3, Vitamin B12), Canola Oil, Artificial Flavor, Cellulose Gel, Soy Lecithin, Tuna Oil, Inositol, Taurine, Cellulose Gum, Potassium Hydroxide, Acesulfame Potassium, Sucralose, L-Carnitine, and Lutein.

Allergens: Contains milk and soy ingredients.

Availability

List Number	Item
67417	PediaSure Peptide 1.5 Cal Vanilla / 8 fl oz (237 mL) Bottle / 24 ct
67419	PediaSure Peptide 1.5 Cal Vanilla / 1000 mL (1.1 QT) Ready-To-Hang Prefilled Container / 8 ct

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Abbott Nutrition Abbott Laboratories
Columbus, OH 43219-3034
1-800-227-5767

2 of 5



PediaSure® Peptide 1.5 Cal

Peptide-Based Nutrition

Nutrition Information - Liquid Vanilla

	8 fl oz (237 mL)		1 L (33.8 fl oz)	
	Value	%RDI / %DV	Value	%RDI / %DV
Protein, g	10.7		45	
Fat, g	14.4		61	
Saturated Fat, g	8		35	
Trans Fat, g	0		0	
Polyunsaturated Fat, g	1.0		4	
Monounsaturated Fat, g	2.5		11	
Cholesterol, mg	15		65	
Carbohydrate, g	47.6		201	
Dietary Fiber, g	1*		5*	
Total Sugars, g	2		7	
Added Sugars, g	2		7	
L-Carnitine, mg	6.0		25	
Taurine, mg	45		190	
Water, g	182		768	
Calories	356		1500	
Vitamin A, mcg RAE	200		850	
Vitamin D, mcg	5.9		25.0	
Vitamin E, mg	3.7		15.5	
Vitamin K, mcg	24		101	
Vitamin C, mg	36		152	
Folate, mcg DFE	180		750	
Folic Acid, mcg	107		450	
Thiamin (Vitamin B1), mg	1.0		4.2	
Riboflavin (Vitamin B2), mg	0.80		3.4	
Vitamin B6, mg	1.0		4.2	
Vitamin B12, mcg	2.1		8.9	
Niacin, mg NE	7.6		31.9	
Choline, mg	120		510	
Biotin, mcg	30		127	
Pantothenic Acid, mg	3.6		15	
Inositol, mg	47		200	
Sodium, mg	255		1075	
Potassium, mg	710		2975	
Chloride, mg	360		1520	
Calcium, mg	500		2090	
Phosphorus, mg	380		1580	
Magnesium, mg	60		255	
Iodine, mcg	35		148	
Manganese, mg	0.7		2.9	

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Nutrition Information - Liquid Vanilla

	8 fl oz (237 mL)		1 L (33.8 fl oz)	
	Value	%RDI / %DV	Value	%RDI / %DV
Copper, mg	0.2		0.89	
Zinc, mg	4.3		18.1	
Iron, mg	5.0		21.1	
Selenium, mcg	12		51	
Chromium, mcg	14		57	
Molybdenum, mcg	14		57	

Liquid Vanilla Footnotes & References

Per 8 fl oz (237 mL)

* 1.1 g from short-chain fructooligosaccharides.

Per 1 L (33.8 fl oz)

* 4.6 g from short-chain fructooligosaccharides.

Preparation

Instructions for Use:

8 fl oz Bottle

- 1 **Shake very well.**
- 1 Store unopened at room temperature.
- 1 After opening, refrigerate and **use within 48 hours.**
- 1 Tube Feeding
 - 1 Follow physician's instructions.
 - 1 Adjust flow rate and volume according to child's condition and tolerance.
 - 1 Additional fluid may be required.
- 1 Oral Feeding
 - 1 Best served chilled.

Ready-To-Hang Container

All medical foods, regardless of type of administration system, require careful handling because they can support microbial growth. Follow these instructions for clean technique and proper setup to reduce the potential for microbial contamination. NOTE: Failure to follow the **Instructions for Use** increases the potential for microbial contamination and may reduce hangtime.

- 1 Administer product at room temperature.
- 1 THOROUGHLY wash hands with soap and water before handling container or feeding set.
- 1 Turn container upside down and SHAKE VIGOROUSLY, using a twisting motion for at least 10 seconds.
- 1 DO NOT touch any part of the container or feeding set that comes into contact with the formula.
- 1 When initiating feeding, follow physician's instructions. Adjust flow rate and volume according to patient's condition and tolerance.
- 1 Additional fluid requirements should be met by giving water between or after feedings or when flushing the tube.
- 1 For Use with Enteral Feeding Pumps:
 - 1 Remove dust cover from Safety Screw Cap.
 - 1 Remove dust cover from feeding set connector.
 - 1 Insert feeding set connector into port of RTH Safety Screw Cap and completely pierce foil.
 - 1 Turn connector collar clockwise until it is securely fastened.
 - 1 Close clamp on feeding set before inverting container.
 - 1 Invert container and suspend, using hanging ring on bottom of container.
 - 1 Unless a shorter hang time is specified by the set manufacturer, hang product for up to 48 hours after initial connection when clean technique and only one new set are used. Otherwise hang for no more than 24 hours.
- 1 Precautions:
 - 1 Follow directions for use provided by manufacturer of feeding sets.
 - 1 Unless a shorter hang time is specified by the set manufacturer, hang product for up to 48 hours after initial connection when clean technique and only one new set are used. Otherwise hang for no more than 24 hours.