

Product Information: Pedialyte® AdvancedCare® Plus

For more information, contact your Abbott Nutrition sales representative or visit
www.abbottnutrition.ca

Pedialyte® AdvancedCare® Plus

Oral rehydration solution for all ages

Usage

- 1 Pedialyte® AdvancedCare® Plus is formulated with 33% more electrolytes* to replenish fluids and electrolytes lost during diarrhea and vomiting to help prevent dehydration. Can be used for maintenance of water and electrolytes following corrective parenteral therapy for diarrhea

Features

- 1 #1 doctor and pharmacist recommended brand.
- 1 33% more electrolytes* for higher electrolyte loss.
- 1 Contains PreActiv® prebiotics in the form of galactooligosaccharides (GOS) that help stimulate the growth of good bacteria in the large intestine.
- 1 Contains the optimal balance of sodium and sugar recommended by experts^{1,2} to help prevent dehydration due to diarrhea and vomiting.
- 1 Helps replenish zinc lost during diarrhea.^{3,4}
- 1 Promotes fluid absorption more effectively than common household beverages.
- 1 1-L reclosable bottles for easy measuring and pouring.
- 1 2 delicious flavours to encourage compliance.
- 1 Low osmolality.
- 1 Gluten-free.
- 1 Kosher.
- 1 Halal.



Safety Precautions

- 1 Not for parenteral use.
- 1 Not for children with milk allergy, GOS allergy or galactosemia.
- 1 For infants under 1 year, consult a physician or qualified health care professional before using Pedialyte® AdvancedCare® Plus as infants and young children are at greater risk of dehydration.

* Sodium electrolytes per equal volume compared to other Pedialyte® products.

* Sodium electrolytes per equal volume compared to other Pedialyte® products.

¹ World Health Organization. Oral Rehydration Salts: Production of the new ORS. 2006.

² Leung A, Prince T. Paediatr Child Health. 2006;11(8):527-31.

³ Bhatnagar S, et al. JPGN. 2004;38(1):34-40

⁴ Lukacik M, et al. Pediatrics. 2008;121(2):326-36.

Ingredients

Liquid Berry Frost:

MEDICINAL INGREDIENTS: Dextrose, Galactooligosaccharides (GOS), Sodium Chloride, Citric Acid, Potassium Citrate, Sodium Citrate, Zinc Gluconate. NON-MEDICINAL INGREDIENTS: Water, Natural and Artificial Flavour, Sucralose, Acesulfame-Potassium, and FD&C Blue #1.

Availability

List Number	Item
6729313	Pedialyte® AdvancedCare® Plus, Berry Frost; 1-L recloseable plastic bottles; 8/case
6729413	Pedialyte® AdvancedCare® Plus, Cherry Pomegranate; 1-L recloseable plastic bottles; 8/case

For more information, contact your Abbott Nutrition sales representative or visit www.abbottnutrition.ca

Abbott Nutrition Abbott Laboratories Co.
Saint-Laurent, Quebec, Canada H4S 1Z6
1-866-767-7411



Nutrition Information - Liquid Berry Frost

	100 mL	1000 mL (1 L)
	Value	Value
ENERGY, Cal (kJ)	10 (42)	100 (418)
MEDICINAL INGREDIENT:		
Sodium, mg (mmol)	138 (6.0)	1380 (60)
Potassium, mg (mmol)	78.0 (2.0)	780 (20)
Chloride, mg (mmol)	177 (5.0)	1770 (50)
Citrate, g (mmol)	0.47 (2.48)	4.7 (24.8)
Zinc, mg	0.78	7.8
Galactooligosaccharides (GOS), g	0.32	3.2
Dextrose, g	1.6	16

Product Characteristics

	Value
Caloric Density (Cal/mL)	0.10
Protein (% of total energy)	0
Protein Source	N/A
Fat (% of total energy)	0
MCT oil (% of total fats)	N/A
Omega-6:Omega-3 Ratio	N/A
Fat Source	N/A
Carbohydrate (% of total energy)	100
Carbohydrate Source	N/A
Total Cal:g Nitrogen Ratio	N/A
Nonprotein Cal:g Nitrogen Ratio	N/A
Water (g/L)	970
Osmolality (mOsm/kg water)	265
Renal Solute Load (mOsm/L)	130
Minimum Tube Size for Gravity/Pump Feeding (Fr)	N/A

For more information, contact your Abbott Nutrition sales representative or visit www.abbottnutrition.ca

Preparation

Directions for Use:

- 1 Tastes best served cold.
- 1 Do not use if tamper band around cap or inner foil seal is missing or broken.
- 1 Remove tamper band. Twist off cap. Remove foil seal.
- 1 Do not add water or dilute.

Recommended dose*

Infants less than 1 year

- 1 Consult a physician before using Pedialyte® AdvancedCare® Plus.

Children 1 year of age

- 1 Give 50-100 mL per episode of diarrhea and/or vomiting, up to a maximum of 1 L over a period of 24 hours.

Children 2–9 years

- 1 Give 100-200 mL per episode of diarrhea and/or vomiting, up to a maximum of 1 L over a period of 8 to 24 hours.

Adults, adolescents and children 10 or more years

- 1 Take 200-400 mL per episode of diarrhea and/or vomiting, or as much as wanted.

*Attempt to give/take solution as soon as diarrhea begins. Take/give the solution frequently and in small amounts (e.g., 5 mL) gradually increased, as tolerated. If more solution is requested/desired, give/take more, as tolerated.

Storage & Handling

Directions for Use:

- 1 After opening, replace cap, refrigerate, and use within 96 hours.
- 1 Store unopened bottles in a cool place. Avoid excessive heat.
- 1 Do not reuse bottles.