

## Product Information: Pedialyte® AdvancedCare®

For more information, contact your Abbott Nutrition sales representative or visit  
[www.abbottnutrition.ca](http://www.abbottnutrition.ca)

# Pedialyte® AdvancedCare®

Oral rehydration solution for all ages

## Usage

- 1 Pedialyte® AdvancedCare® contains PreActiv® prebiotics and is specially formulated to help prevent dehydration in children and adults by replenishing fluids and electrolytes lost during diarrhea and vomiting. Can be used for maintenance of water and electrolytes following corrective parenteral therapy for diarrhea.

## Features

- 1 #1 doctor and pharmacist recommended brand.
- 1 Contains PreActiv® prebiotics in the form of galactooligosaccharides (GOS) that help stimulate the growth of good bacteria in the large intestine.
- 1 Contains the optimal balance of sodium and sugar recommended by experts<sup>1,2</sup> to help prevent dehydration due to diarrhea and vomiting.
- 1 Helps replenish zinc lost during diarrhea.<sup>3,4</sup>
- 1 Promotes fluid absorption more effectively than common household beverages.
- 1 1-L reclosable bottles for easy measuring and pouring.
- 1 2 delicious flavours to encourage compliance.
- 1 Low osmolality.
- 1 Low-residue.
- 1 Gluten-free.
- 1 Kosher.
- 1 Halal (Blue Raspberry only).



## Safety Precautions

- 1 Not for parenteral use.
- 1 Not for children with milk allergy, GOS allergy or galactosemia.
- 1 For infants under 1 year, consult a physician or qualified health care professional before using Pedialyte® AdvancedCare® as infants and young children are at greater risk of dehydration.

<sup>1</sup> World Health Organization. Oral Rehydration Salts: Production of the new ORS. 2006.

<sup>2</sup> Leung A, Prince T. Paediatr Child Health. 2006;11(8):527-31.

<sup>3</sup> Bhatnagar S, et al. JPGN. 2004;38(1):34-40

<sup>4</sup> Lukacik M, et al. Pediatrics. 2008;121(2):326-36.

## Ingredients

### Liquid Cherry Punch:

MEDICINAL INGREDIENTS: Dextrose, Galactooligosaccharides, Citric Acid, Potassium Citrate, Sodium Chloride, Sodium Citrate, Zinc Gluconate. NON-MEDICINAL INGREDIENTS: Water, Natural and Artificial Flavours, Sucralose, Acesulfame-Potassium, and FD&C Red #40.

## Availability

List Number	Item
6647013	Pedialyte® AdvancedCare®, Cherry Punch; 1-L reclosable plastic bottles; 8/case
6647113	Pedialyte® AdvancedCare®, Blue Raspberry; 1-L reclosable plastic bottles; 8/case

For more information, contact your Abbott Nutrition sales representative or visit [www.abbottnutrition.ca](http://www.abbottnutrition.ca)

Abbott Nutrition Abbott Laboratories Co.  
Saint-Laurent, Quebec, Canada H4S 1Z6  
1-866-767-7411

## Nutrition Information - Liquid Cherry Punch

	1000 mL (1 L)
	Value
ENERGY, Cal (kJ)	100 (418)
MEDICINAL INGREDIENT:	
Sodium, mg (mmol)	1030 (45)
Potassium, mg (mmol)	782 (20)
Chloride, mg (mmol)	1244 (35)
Citrate, g (mmol)	4.8 (25.5)
Zinc, mg	7.8
Galactooligosaccharides (GOS), g	3.2
Dextrose, g	16.0

## Product Characteristics

	Value
Caloric Density (Cal/mL)	0.10
Protein (% of total energy)	0
Protein Source	N/A
Fat (% of total energy)	0
MCT oil (% of total fats)	N/A
Omega-6:Omega-3 Ratio	N/A
Fat Source	N/A
Carbohydrate (% of total energy)	100
Carbohydrate Source	N/A
Total Cal:g Nitrogen Ratio	N/A
Nonprotein Cal:g Nitrogen Ratio	N/A
Water (g/L)	970
Osmolality (mOsm/kg water)	240
Renal Solute Load (mOsm/L)	100
Minimum Tube Size for Gravity/Pump Feeding (Fr)	N/A

For more information, contact your Abbott Nutrition sales representative or visit [www.abbottnutrition.ca](http://www.abbottnutrition.ca)

## Preparation

### Directions for Use:

- 1 Tastes best served cold.
- 1 Do not use if tamper band around cap or inner foil seal is missing or broken.
- 1 To open: Remove tamper band. Twist off cap. Remove foil seal.
- 1 Do not add water or dilute.

### Recommended dose

- 1 Infants less than 1 year of age
  - 1 **Consult a physician before using Pedialyte AdvancedCare.**
- 1 Children 1 years of age\*
  - 1 Give 50-100 mL per episode of diarrhea and/or vomiting, up to a maximum of 1 L over a period of 24 hours.
- 1 Children 2–9 years\*
  - 1 Give 100-200 mL per episode of diarrhea and/or vomiting, up to a maximum of 1 L over a period of 8 to 24 hours.
- 1 Adults, adolescents and children 10 or more years\*
  - 1 Take 200-400 mL per episode of diarrhea and/or vomiting, or as much as wanted.

\* Attempt to give/take solution as soon as diarrhea begins. Take/give the solution frequently and in small amounts (e.g. 5 mL), gradually increased, as tolerated. If more solution is requested/desired, give/take more, as tolerated. For dehydration due to other causes, consult a health care professional

## Storage & Handling

Store unopened bottles in a cool place. Avoid excessive heat. Do not reuse bottle. After opening, replace cap, refrigerate, and use within 96 hours or discard.