

## Product Information: Pedialyte® Classic

For more information, contact your Abbott Nutrition Representative or visit  
[www.abbottnutrition.com](http://www.abbottnutrition.com)

# Pedialyte® Classic

## Electrolyte solution

- 1 PEDIALYTE CLASSIC is an advanced rehydration solution specially formulated with an optimal balance of glucose and electrolytes needed to help replenish vital fluids and minerals which, when lost, can lead to dehydration. It can help prevent mild to moderate dehydration caused by vomiting, diarrhea, exercise, travel, and extreme heat.
- 1 Use under medical supervision for the dietary management of mild to moderate dehydration during diarrhea and vomiting.
- 1 #1 Pediatrician & Pharmacist recommended brand for hydration.
- 1 Medical-grade hydration.
- 1 Designed to promote fluid absorption more effectively than common household beverages.
- 1 Has glucose to promote sodium and water absorption.
- 1 Quickly replaces fluids and electrolytes.
- 1 Unflavored liquid available for infants.
- 1 Ready-to-use reclosable plastic liter bottles allow easy measuring and pouring.
- 1 Low osmolality.
- 1 Contains no fruit juice.
- 1 Suitable for lactose intolerance.
- 1 Gluten-free.
- 1 Kosher.
- 1 Halal.



## Safety Precautions

- 1 No mixing or diluting is necessary or recommended.
- 1 For children under 1 year of age, consult your doctor.
- 1 **Administration Guide for Infants and Young Children**

Age	Approximate Weight <sup>1</sup>		PEDIALYTE for Maintenance <sup>2</sup>
	lb	kg	fl oz/day
Weeks			
2	9	4.0	16 to 20
Months			
3	14	6.4	30 to 34
6	18	8.2	36 to 42
9	21	9.5	39 to 45
Years			
1	23	10.5	42 to 47
1?	26	11.8	47 to 52
2	28	12.7	48 to 53
2?	30	13.6	51 to 56
3	32	14.4	53 to 57
3?	34	15.3	54 to 57
4	36	16.3	55 to 59

Administration Guide does not apply to infants younger than 1 week of age. For children older than 4 years of age, maintenance intakes may exceed 2 liters daily. If there is vomiting or fever, or if diarrhea continues beyond 24 hours, consult the child's physician.

<sup>1</sup> Weight based on the 50th percentile of weight for age for boys from the National Center for Health Statistics (NCHS) Centers for Disease Control and Prevention (CDC) growth charts. Kuczmarski RJ, Ogden CL, Grummer-Strawn LM, et al: CDC Growth Charts: United States. Data from Vital and Health Statistics of the Centers of Disease Control and

For more information, contact your Abbott Nutrition Representative or visit [www.abbottnutrition.com](http://www.abbottnutrition.com)

Prevention/National Center for Health Statistics. *Advance Data*, no. 314, December 4, 2000.

<sup>2</sup> Fluid intake is total fluid requirement from oral electrolyte solution, formula, or other fluids, but does not take into account ongoing stool losses. Fluid loss in the stool should be replaced by consumption of an extra amount of Pedialyte equal to stool losses, in addition to the fluid maintenance requirement in the Administration Guide.

### 1 Dosage

Refer to Administration Guide to restore fluid and minerals lost in diarrhea and vomiting. Pedialyte should be offered frequently in amounts tolerated. Total daily intake should be adjusted to meet individual needs, based on thirst and response to therapy. The suggested intakes for maintenance are based on water requirements for ordinary energy expenditure. (Extrapolated from Barnes LA, Curran JS: Nutrition, in Nelson WE (sr ed), Behrman RE, Kliegman RM, Arvin AM (eds): *Nelson Textbook of Pediatrics*, ed 15. Philadelphia: WB Saunders Co, 1996, pp 141-143.)

## Ingredients

### Liquid Grape:

Water, Dextrose. Less than 0.5% of: Citric Acid, Potassium Citrate, Salt, Natural and Artificial Flavor, Sodium Citrate, Sucralose, Acesulfame Potassium, Zinc Gluconate, Red 40, and Blue 1.

## Availability

List Number	Item
00240	Pedialyte Grape / 33.8 fl oz (1 L) Bottle / 2 x 8 ct
00336	Pedialyte Unflavored / 33.8 fl oz (1 L) Bottle / 2 x 4 ct
00365	Pedialyte Mixed Fruit / 33.8 fl oz (1 L) Bottle / 2 x 8 ct
53983	Pedialyte Strawberry / 33.8 fl oz (1 L) Bottle / 8 ct
64380	Pedialyte Unflavored / 33.8 fl oz (1 L) Bottle / 4 ct
66284	Pedialyte Mixed Fruit / 33.8 fl oz (1 L) Bottle / 4 ct
67461	Pedialyte Coconut Burst / 33.8 fl oz (1 L) Bottle / 4 ct
69090	Cherry Frost / 1.1 QT (1 L) Bottle / 4 ct
69096	Pedialyte Fruit Punch / 1.1 QT (1 L) Bottle / 4 ct
69098	Pedialyte Iced Berry / 1.1 QT (1 L) Bottle / 4 ct

For more information, contact your Abbott Nutrition Representative or visit  
[www.abbottnutrition.com](http://www.abbottnutrition.com)

## Nutrition Information - Liquid Grape

	12 fl oz (360 mL)		33.8 fl oz (1 L)	
	Value	%RDI / %DV*	Value	%RDI / %DV*
Calories	25		70	
Total Fat, g	0	0	0	0
Sodium, mg	390	17	1080	47
Total Carbohydrate, g	7	3	19	7
Total Sugars, g	5		13	
Added Sugars, g	5	10	13	26
Protein, g	0		0	
Potassium, mg	280	6	780	15
Zinc, mg	2.8	25	7.8	70
Chloride, mg	440	20	1240	50

## Liquid Grape Footnotes & References

### Per 12 fl oz (360 mL)

\*The % Daily Value (DV) tells you how much a nutrient in a serving of food contributes to a daily diet. 2,000 calories a day is used for general nutrition advice. -- Not a significant source of saturated fat, trans fat, cholesterol, dietary fiber, vitamin D, calcium and iron.

### Per 33.8 fl oz (1 L)

\*The % Daily Value (DV) tells you how much a nutrient in a serving of food contributes to a daily diet. 2,000 calories a day is used for general nutrition advice. -- Not a significant source of saturated fat, trans fat, cholesterol, dietary fiber, vitamin D, calcium and iron.

## Preparation

### Instructions for Use:

- 1 Use under medical supervision.
- 1 READY TO USE. DO NOT ADD WATER OR DILUTE.
- 1 Tastes best served cold.
- 1 For children under 1 year of age: consult your doctor.
- 1 To rehydrate after fluid loss: begin with small frequent sips. 1-2 liters per day may be needed while symptoms persist. Consult your doctor if vomiting, fever, or diarrhea continues beyond 24 hours or if consumption needs are greater than 2 liters per day.
- 1 For general hydration, sip as needed.

### Storage Instructions:

- 1 Do not use if printed band around cap or inner foil seal is missing or broken.
- 1 Remove protective band and twist off cap.
- 1 Remove and throw away protective foil seal.
- 1 After opening, replace cap, refrigerate and use within 48 hours.
- 1 Store bottles upright in a cool place. Avoid excessive heat.
- 1 Do not reuse bottle.