

Product Information: ProViMin®

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com



- 1 For use in the management of patients who require a formula modified in carbohydrate, fat, and/or increased protein: abetalipoproteinemia; cholestasis; chylothorax; fatty acid oxidation defects; glutaric aciduria type II; hyperlipoproteinemia type I (fasting chylomicronemia); hypobetalipoproteinemia; lymphangiectasis, intestinal malabsorption of carbohydrate and/or fat; supplement for any patient who requires increased protein, minerals, and vitamins; X-linked adrenoleukodystrophy.
- 1 Use under medical supervision.
- 1 High-quality source of protein (casein).
- 1 Virtually carbohydrate- and fat-free; provides flexibility in prescribing a formula in which the amount and type of carbohydrate and fat can be added per individual infant's need and tolerance.
- 1 Meets the Dietary Reference Intakes and the American Academy of Pediatrics Committee on Nutrition (AAPCON) recommendations for vitamin and mineral intakes for infants when fed at a dilution of 3.25 g protein/100 Cal.
- 1 Calcium-to-phosphorus ratio of 1.4:1, which is similar to that of standard infant formulas and meets AAPCON recommendations for that ratio.
- 1 L-carnitine (40 mg/100 g) and taurine (110 mg/100 g).
- 1 Lactose-free.
- 1 Gluten-free.
- 1 Halal.
- 1 Kosher.

Safety Precautions

- 1 Does not supply sufficient amounts of energy as carbohydrate, fat or linoleic acid. These nutrients should be supplied from other sources under the supervision of a physician.
- 1 Not for IV use.
- 1 Do not boil mixture or use terminal sterilization.
- 1 **Never use a microwave oven to prepare or warm mixture.** Serious burns can result.

Ingredients

Powder Unflavored:

Sodium Caseinate, Calcium Phosphate, Potassium Citrate, Potassium Chloride. Less than 2% of: Magnesium Chloride, Ascorbic Acid, Coconut Oil, Choline Chloride, Ferrous Sulfate, Taurine, m-Inositol, L-Methionine, d-Alpha-Tocopheryl Acetate, Zinc Sulfate, L-Carnitine, Niacinamide, Calcium Pantothenate, Vitamin A Palmitate, Copper Sulfate, Thiamine Hydrochloride, Riboflavin, Pyridoxine Hydrochloride, Folic Acid, Manganese Sulfate, Potassium Iodide, Phylloquinone, Biotin, Sodium Selenate, Vitamin D3, Vitamin B12, Salt, and Calcium Carbonate.

Allergens: Contains milk ingredients.

Availability

List Number	Item
50260	ProViMin Powder / 5.3 oz (150 g) Can / 6 ct

Nutrition Information - Powder Unflavored

	100 g Powder		100 Cal (5 fl oz)	
	Value	%RDI / %DV	Value	%RDI / %DV
Protein, g	73		23	
Fat, g	1.4		0.4	
Carbohydrate, g	2.0		0.6	
Water, g	4.0			
Linoleic Acid, mg	0.0		0.0	
Potential Renal Solute Load, mOsm	673		215	
Calories	313		100	
Vitamin A, IU	6740		2150	
Vitamin D, IU	1000		320	
Vitamin E, IU	67		21	
Vitamin K, mcg	90		29	
Thiamin (Vitamin B1), mcg	2240		716	
Riboflavin (Vitamin B2), mcg	2020		645	
Vitamin B6, mcg	1350		431	
Vitamin B12, mcg	5.6		1.8	
Niacin, mcg	24000		7670	
Folic Acid (Folacin), mcg	320		102	
Pantothenic Acid, mcg	10100		3230	
Biotin, mcg	100		32	
Vitamin C (Ascorbic Acid), mg	200		64	
Choline, mg	335		107	
Inositol, mg	105		34	
Calcium, mg	2400		767	
Calcium, mEq	120		38.4	
Phosphorus, mg	1700		543	
Magnesium, mg	200		64	
Iron, mg	40		13	
Zinc, mg	17		5.4	
Manganese, mcg	200		64	
Copper, mcg	2100		671	
Iodine, mcg	335		107	
Selenium, mcg	46		13	
Sodium, mg	1200		383	
Sodium, mEq	52		17	
Potassium, mg	3300		1050	
Potassium, mEq	84		27	
Chloride, mg	2300		735	
Chloride, mEq	65		21	

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Preparation

Directions for Preparation and Use: Use Only As Directed by a Physician

Depending on specific needs of each individual, preparation will vary as prescribed by physician. Failure to follow these directions could result in harm. If feeding an infant, ask your baby's doctor if you need to use cooled, boiled water for mixing and boil (sterilize) bottles, nipples and rings before use.

- 1 Wash your hands, surfaces and utensils.
- 1 Pour a small amount of the prescribed liquid into a clean container or blender.
- 1 Add the prescribed amount of ProViMin powder and mix or blend on low speed to make a smooth paste.
- 1 Gradually add remaining liquid (and other ingredients if recommended) to the mixture while stirring or blending. Stir or blend until well blended.
- 1 Pour mixture into clean feeding container; cap.
- 1 Shake well before feeding.

Once feeding begins, **use within 1 hour or discard.**

Approximate Weights for Unpacked, Level U.S. Standard Dry Household Measures for Powder*

Household Measure (US)	Approx. Weight (g)
1 Tbsp	2.9
1/4 Cup	11.0
1/2 Cup	20.0
2/3 Cup	30.0
1 Cup	44.0

* For most accurate results, ProViMin should be weighed on a scale that reads in grams.

Approximate Weights for Unpacked, Level U.S. Standard Dry Household Measures for Powder*

Household Measure (US)	Approx. Weight (g)
1 Tbsp	2.9
1/4 Cup	11.0
1/2 Cup	20.0
2/3 Cup	30.0
1 Cup	44.0

* For most accurate results, ProViMin should be weighed on a scale that reads in grams.

Storage & Handling

Storage

- 1 Once mixed, store container in refrigerator and **use within 24 hours.**
- 1 Store unopened or opened can at room temperature; avoid extreme temperatures.
- 1 **Use opened can contents within 1 month.**