Product Information: Propimex®-1

Propimex®-1

Amino acid-modified infant formula with iron

- PROPIMEX-1 is for nutrition support of infants and toddlers with propionic or methylmalonic acidemia.
- Use under medical supervision.
- Methionine- and valine-free to allow greater intake of intact protein; low in isoleucine and threonine.
- 1 L-carnitine (900 mg/100 g) to help supply amount normally found in human milk and foods of animal origin and to help excrete toxic metabolites.
- 1 Taurine (40 mg/100 g) to help supply amount normally found in human milk and foods of animal origin.
- Approximately 40% of energy as fat to help achieve acceptable formula osmolality.
- 1 6.6% of energy as linoleic acid.
- 1 Nutrient profile specifically designed for infants and toddlers.
- 1 When fed according to the Abbott Nutrition Support Protocols, provides adequate amounts of all nutrients.
- 1 Non-GMO.
- 1 DHA and ARA to help support brain and eye development.
- 1 Lactose-free.
- Gluten-free.
- 1 Kosher made with pareve ingredients.
- Halal.
- Includes DHA/ARA.

Safety Precautions

- Not intended as a sole source of nutrition.
- 1 Must be supplemented with protein and fluid in prescribed amounts to completely meet isoleucine, methionine, threonine, valine and water requirements.
- 1 Do not boil mixture or use terminal sterilization.
- 1 Never use a microwave oven to warm formula. Serious burns can result.
- 1 Not for IV use.
- 1 Powdered infant formulas are not sterile and should not be fed to premature infants or infants who might have immune problems unless directed and supervised by your baby's doctor.

Ingredients

Powder Unflavored:

Corn Syrup Solids, High Oleic Safflower Oil, Coconut Oil, Soy Oil, L-Alanine, L-Glutamic Acid, Modified Corn Starch. Less than 2% of: L-Proline, L-Aspartic Acid, L-Arginine, L-Lysine, L-Leucine, Calcium Phosphate, Potassium Phosphate, L-Carnitine, L-Tyrosine, L-Phenylalanine, L-Serine, Sodium Citrate, L-Cystine Dihydrochloride, Potassium Citrate, Glycine, Magnesium Chloride, L-Histidine, M. Alpina Oil^{*}, Schizochytrium Sp. Oil[†], Calcium Carbonate, Citric Acid, Ascorbic Acid, L-Tryptophan, Inositol, Choline Chloride, L-Isoleucine, L-Threonine, Taurine, Ferrous Sulfate, Ascorbyl Palmitate, Zinc Sulfate, dl-Alpha-Tocopheryl Acetate, Mixed Tocopherols, Calcium Pantothenate, Niacinamide, Manganese Sulfate, Thiamine Hydrochloride, Copper Sulfate, Vitamin A Palmitate, Riboflavin, Pyridoxine Hydrochloride, Potassium Iodide, Folic Acid, Phylloquinone, Chromium Chloride, Sodium Selenate, Sodium Molybdate, Biotin, Vitamin D3, Vitamin B12, and Salt.

Availability

List Number	Item
67058	Propimex-1 / 14.1 oz (400 g) Can / 6 ct







Ingredients not genetically engineered.

[†] Severe burn injuries, trauma, post-op recovery.

Source of ARA.

[†] Source of DHA.

Nutrition Information - Powder Unflavored

	ored 100 g	100 g Powder	
	Value	%RDI / %DV	
		/0KDI / /0DV	
Protein Equivalent, g	15		
Fat, g	21.7		
Carbohydrate, g	53		
Linoleic Acid, mg	3650		
Linolenic Acid, mg	360		
L-Carnitine, mg	900		
Calories	480		
Vitamin A, IU	1300		
Vitamin A, mcg RE	390		
Vitamin D, IU	450		
Vitamin D, mcg	11.3		
Vitamin E, IU	12.5		
Vitamin E, mg alpha-TE	6.4		
Vitamin K, mcg	60		
Thiamin (Vitamin B1), mcg	1120		
Riboflavin (Vitamin B2), mcg	650		
Vitamin B6, mcg	560		
Vitamin B12, mcg	0.85		
Niacin, mcg	4450		
Niacin, mg NE	7.7		
Folic Acid (Folacin), mcg	88		
Pantothenic Acid, mcg	4000		
Biotin, mcg	23.5		
Vitamin C, mg	75		
Choline, mg	95		
Inositol, mg	144		
Calcium, mg	575		
Phosphorus, mg	400		
Magnesium, mg	50		
Iron, mg	7.6		
Zinc, mg	5.2		
Manganese, mcg	385		
Copper, mcg	408		
lodine, mcg	65		
Selenium, mcg	21		
Chromium, mcg	12		
Molybdenum, mcg	13.8		
Sodium, mg	190		
Sodium, mEq	8.3		

For more information, contact your Abbott Nutrition Representative or visit www.abbottnutrition.com



Nutrition Information - Powder Unflavored

	100 g Powder	
	Value	%RDI / %DV
Potassium, mg	675	
Potassium, mEq	17.3	
Chloride, mg	410	
Chloride, mEq	11.6	
Potential Renal Solute Load, mOsm	136	

Preparation

Instructions for Use: Use Only As Directed by a Physician

Your baby's health depends on carefully following these directions. Proper hygiene, handling and storage are important when preparing infant formula. Failure to follow these directions could result in severe harm. Ask your baby's doctor if you need to use cooled, boiled water for mixing and if you need to boil (sterilize) bottles, nipples and rings before use.

- 1 Wash your hands, surfaces and utensils.
- 1 Pour prescribed amount of water into clean container.
- 1 Add prescribed amount of Propimex-1 powder (and other ingredients if recommended).
- 1 Shake well for 10-15 seconds; if using a blender, mix no more than 5 seconds.
- 1 Pour mixture into clean feeding bottles or container; cap.
- Once feeding begins, use within 1 hour or discard.

Approximate weights for upacked, level U.S. standard dry household measures for powder*

Household Measure (US)	Weight (g)	
1 Tbsp	8	
1/4 Cup	30	
1/3 Cup	40	
1/2 Cup	60	
1 Cup	120	

^{*} For most accurate results, Propimex-1 should be weighed on a scale that reads in grams.

Storage Instructions:

- 1 Once mixed, store bottles or container in refrigerator and feed to baby or child within 24 hours
- Store unopened or opened can at room temperature; avoid extreme temperatures.
- 1 Use opened can contents within 1 month.

