

Product Information: Pulmocare®

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

- 1 PULMOCARE is designed for people with chronic obstructive pulmonary disease (COPD), cystic fibrosis, or respiratory failure who may benefit from a high-calorie, modified carbohydrate and fat enteral formula that may help reduce diet-induced carbon dioxide production.
- 1 For tube or oral feeding.
- 1 For supplemental or sole-source nutrition.
- 1 Use under medical supervision.
- 1 Provides complete and balanced nutrition.
- 1 20% of fat as MCTs to enhance fat absorption.
- 1 Elevated antioxidants vitamins C & E, and beta-carotene.
- 1 Kosher.
- 1 Halal.
- 1 Gluten-free.
- 1 Suitable for lactose intolerance.



Safety Precautions

- 1 Not for IV use.
- 1 Not suitable for people with galactosemia.

Ingredients

Liquid Vanilla:

Water, Sodium Caseinate, Sugar, Canola Oil, Corn Maltodextrin, Medium Chain Triglycerides, Corn Oil, Calcium Caseinate, High Oleic Safflower Oil. Less than 0.5% of: Natural & Artificial Flavors, Magnesium Chloride, Calcium Phosphate, Soy Lecithin, Potassium Citrate, Sodium Citrate, Potassium Phosphate, Sodium Hydroxide, Choline Chloride, Ascorbic Acid, L-Carnitine, Taurine, Gellan Gum, Salt, Zinc Sulfate, d-Alpha-Tocopheryl Acetate, Ferrous Sulfate, Niacinamide, Calcium Pantothenate, Manganese Sulfate, Copper Sulfate, Thiamine Hydrochloride, Pyridoxine Hydrochloride, Riboflavin, Beta-Carotene, Vitamin A Palmitate, Folic Acid, Biotin, Chromium Chloride, Sodium Molybdate, Potassium Iodide, Sodium Selenate, Phylloquinone, Vitamin B12, and Vitamin D3.

Allergens: Contains milk and soy ingredients.

Availability

List Number	Item
62725	Pulmocare / 33.8 fl oz (1 L) Ready-To-Hang Prefilled Container / 8 ct
64811	Pulmocare Vanilla / 8 fl oz (237 mL) Recloseable Carton / 24 ct

Nutrition Information - Liquid Vanilla

	8 fl oz (237 mL)	
	Value	%RDI / %DV
Protein, g	14.8	
Fat, g	22.1	
Carbohydrate, g	25	
Total Sugars, g	14	
Added Sugars, g	14	
L-Carnitine, mg	36	
Taurine, mg	36	
Water, g	186	
Calories	355	
Vitamin A, mcg RAE	852*	
Vitamin D, mcg	2.5	
Vitamin E, mg	15.8	
Vitamin K, mcg	20	
Vitamin C, mg	75	
Folate, mcg DFE	333	
Folic Acid, mcg	200	
Thiamin (Vitamin B1), mg	0.75	
Riboflavin (Vitamin B2), mg	0.85	
Vitamin B6, mg	1	
Vitamin B12, mcg	3	
Niacin, mg NE	11.1	
Choline, mg	125	
Biotin, mcg	150	
Pantothenic Acid, mg	5	
Sodium, mg	310	
Potassium, mg	422	
Chloride, mg	400	
Calcium, mg	250	
Phosphorus, mg	250	
Magnesium, mg	100	
Iodine, mcg	38	
Manganese, mg	1.3	
Copper, mg	0.5	
Zinc, mg	5.7	
Iron, mg	4.5	
Selenium, mcg	18	
Chromium, mcg	30	
Molybdenum, mcg	38	

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Liquid Vanilla Footnotes & References

Per 8 fl oz (237 mL)

*477 mcg RAE of vitamin A activity supplied by 1.2 mg of beta-carotene.

Preparation

Instructions for Use:

8 fl oz recloseable container

- 1 Store unopened at room temperature.
- 1 Shake well prior to opening.
- 1 Once opened, reclose, refrigerate and **use within 48 hours**.

Oral Feeding:

- 1 May be used for total or supplemental nutrition.
- 1 May be fed at room temperature or chilled.

Tube Feeding:

- 1 Follow physician's instructions.
- 1 Adjust flow rate and volume according to patient's condition and tolerance.
- 1 Feed at room temperature using a feeding pump or syringe.
- 1 Additional fluid requirements should be met by giving water between or after feedings or when flushing the tube.
- 1 Avoid contamination during preparation and use.

Ready-To-Hang Container

All medical foods, regardless of type of administration system, require careful handling because they can support microbial growth. Follow these instructions for clean technique and proper setup to reduce the potential for microbial contamination.

NOTE: Failure to follow the Instructions for Use increases the potential for microbial contamination and reduces hangtime.

- 1 Administer product at room temperature.
- 1 THOROUGHLY wash hands with soap and water before handling container or feeding set.
- 1 Turn container upside down and SHAKE VIGOROUSLY, using a twisting motion for at least 10 seconds.
- 1 DO NOT touch any part of the container or feeding set that comes into contact with the formula.
- 1 When initiating feeding, follow physician's instructions. Adjust flow rate and volume according to patient's condition and tolerance.
- 1 Additional fluid requirements should be met by giving water between or after feedings or when flushing the tube.

For Use with Enteral Feeding Pumps:

- 1 Remove dust cover from RTH Safety Screw Cap.
- 1 Remove dust cover from feeding set connector.
- 1 Insert feeding set connector into port of RTH Safety Screw Cap and completely pierce foil.
- 1 Turn connector collar clockwise until it is securely fastened.
- 1 Close clamp on set before inverting container.
- 1 Invert container and suspend, using hanging ring on bottom of container.

Precautions:

- 1 Follow directions for use provided by manufacturer of feeding sets.
- 1 Unless a shorter hang time is specified by the set manufacturer, hang product for up to 48 hours after initial connection when clean technique and only one new set are used. Otherwise hang for no more than 24 hours.