Milk-based infant formula with iron

# **Product Information:** Similac® 360 Total Care®

#### Milk-based infant formula with iron

- SIMILAC 360 TOTAL CARE is a 20 Cal/fl oz, nutritionally complete, non-GMO, milk-based, iron fortified infant formula designed to support the immune system, brain development and digestive health. Key ingredients found in breast milk including 5 HMO prebiotics. For use as a supplement or alternative to breastfeeding.
- Designed to support the immune system, brain development, and digestive health.
- 5 HMO Prebiotics. HMOs are immune nourishing prebiotics structurally identical to those in breast milk.<sup>1</sup>
- 1 DHA, lutein, vitamin E.
- 1 Similac 360 Total Care is our closest formula to breast milk.
- 1 Similac 360 Total Care may soften stools to be more like those of the breastfed infant.
- No artificial growth hormones.<sup>†</sup>
- 1 Non-GMO.\*
- 1 No palm olein oil.
- 1 Halal.
- 1 Kosher made with dairy ingredients.

### **Safety Precautions**

- Not for infants or children with galactosemia.
- 1 Powdered infant formulas are not sterile and should not be fed to premature infants or infants who might have immune problems unless directed and supervised by your baby's doctor.
- 1 Never use a microwave oven to warm formula. Serious burns can result.

## **Ingredients**

#### **Powder Unflavored:**

Nonfat Milk, Lactose, High Oleic Safflower Oil, Whey Protein Concentrate, Soy Oil, Coconut Oil, Human Milk Oligosaccharides (2'-Fucosyllactose, Lacto-N-Tetraose, 3-Fucosyllactose, 6'-Sialyllactose, 3'-Sialyllactose)\*. Less than 2% of: Schizochytrium Sp. Oil<sup>†</sup>, M. Alpina Oil<sup>‡</sup>, Beta-Carotene, Lutein, Potassium Citrate, Calcium Carbonate, Ascorbic Acid, Soy Lecithin, Potassium Chloride, Magnesium Chloride, Ferrous Sulfate, Choline Bitartrate, Choline Chloride, Ascorbyl Palmitate, Salt, Taurine, Inositol, Zinc Sulfate, Mixed Tocopherols, d-Alpha-Tocopheryl Acetate, Niacinamide, Calcium Pantothenate, L-Carnitine, Vitamin A Palmitate, Copper Sulfate, Thiamine Hydrochloride, Riboflavin, Pyridoxine Hydrochloride, Folic Acid, Manganese Sulfate, Phylloquinone, Biotin, Sodium Selenate, Vitamin D3, Vitamin B12, Calcium Phosphate, Potassium Phosphate, Potassium Iodide, Potassium Hydroxide, and Nucleotides (Adenosine 5'-Monophosphate, Cytidine 5'-Monophosphate, Disodium Guanosine 5'-Monophosphate).

Allergens: Contains milk and soy ingredients.





<sup>\*</sup> Ingredients not genetically engineered.

<sup>\*</sup> Ingredients not genetically engineered.

<sup>&</sup>lt;sup>†</sup> Milk ingredients are derived from milk of non-rbST-treated cows, which is not different from milk of rbST-treated cows.

<sup>&</sup>lt;sup>1</sup> Hill DR, Chow JM, Buck RH. Nutrients. 2021;13(10):3364.

<sup>\*</sup> HMO, not from human milk.

<sup>&</sup>lt;sup>†</sup> Source of DHA.

<sup>&</sup>lt;sup>‡</sup> Source of ARA.

Milk-based infant formula with iron

## **Availability**

List Number	Item
68063	Similac 360 Total Care Powder / 20.6 oz (584 g) / 4 ct
68065	Similac 360 Total Care Powder / 30.3 oz (859 g) / 4 ct
68067	Similac 360 Total Care Powder / 40 oz (1.13 kg) / 6 ct
68111	Similac 360 Total Care Powder / 36 oz (1.02 kg) Can /6 ct
68113	Similac 360 Total Care Powder / 0.60 oz (17.2 g) Packets / 4 x 16 ct
68116	Similac 360 Total Care Powder / 30.8 oz (873 g) Can / 4 ct
68130	Similac 360 Total Care Ready to Feed / 2 fl oz (59 mL) Bottle with Nipple / 4 x 6 ct
68133	Similac 360 Total Care Ready to Feed / 2 fl oz (59 mL) Bottle / 2 x 12 ct
68136	Similac 360 Total Care Ready to Feed / 2 fl oz (59 mL) Bottle / 48 ct
68138	Similac 360 Total Care Ready to Feed / 8 fl oz (237 mL) Bottle / 4 x 6 ct
68145	Similac 360 Total Care Ready To Feed / 32 fl oz (946 mL) Bottle / 6 ct
68654	Similac 360 Total Care Powder / 30.8 oz (873 g) Can / 6 ct



Milk-based infant formula with iron

### **Nutrition Information - Powder Unflavored**

Nutrition Information - Powd		1000 ml /1.1 OT propared as			
	100 Cal (5 fl oz, p	100 Cal (5 fl oz, prepared as directed)		directed)	
	Value	%RDI / %DV	Value	%RDI / %DV	
Calories	100		676		
Volume, mL	148		1000		
Protein, g	2.07		14		
Fat, g	5.6		37.87		
Carbohydrate, g	10.5		71.01		
Water, g	134		900		
Linoleic Acid, mg	1000		6763		
Vitamin A, IU	300		2029		
Vitamin D, IU	60		406		
Vitamin E, IU	1.5		10.1		
Vitamin K, mcg	8		54.1		
Thiamin (Vitamin B1), mcg	100		676		
Riboflavin (Vitamin B2), mcg	150		1014		
Vitamin B6, mcg	60		406		
Vitamin B12, mcg	0.25		1.69		
Niacin, mcg	1050		7101		
Folic Acid (Folacin), mcg	15		101.4		
Pantothenic Acid, mcg	450		3043		
Biotin, mcg	4.4		29.8		
Vitamin C (Ascorbic Acid), mg	9		60.9		
Choline, mg	24		162.3		
lnositol, mg	24		162.3		
Calcium, mg	78		528		
Calcium, mEq	3.9		26.4		
Phosphorus, mg	42		284		
Magnesium, mg	6		40.6		
Iron, mg	1.8		12.17		
Zinc, mg	0.75		5.07		
Manganese, mcg	5		33.8		
Copper, mcg	90		609		
lodine, mcg	15		101		
Selenium, mcg	2		14		
Sodium, mg	24		162		
Sodium, mEq	1		6.8		
Potassium, mg	105		710		
Potassium, mEq	2.7		18.3		
Chloride, mg	65		440		
Chloride, mEq	1.8		12.2		

For more information, contact your Abbott Nutrition Representative or visit www.abbottnutrition.com



Milk-based infant formula with iron

Milk-based infant formula with iron

## **Preparation**

#### Instructions for Use:

#### Powder

Your baby's health depends on carefully following these directions. Proper hygiene, handling and storage are important when preparing infant formula. Failure to follow these directions could result in severe harm. Ask your baby's doctor if you need to use cooled, boiled water for mixing and if you need to boil (sterilize) bottles, nipples and rings before use.

#### Can or SimplePac

- 1 Wash your hands, surfaces and utensils.
- 1 Pour water into clean bottle.
- 1 Add 1 unpacked level scoop (8.6 g) to each 2 fl oz of water.
- 1 Return dry scoop to container.
- 1 Cap bottle; shake well; attach nipple.
- 1 Once feeding begins, use within 1 hour or discard.

#### Powder Single

- 1 Wash your hands, surfaces and utensils.
- Pour 4 fl oz of water into clean bottle.
- 1 Slowly add 1 powder single (17.2 g) to bottle.
- 1 Cap bottle; shake well; attach nipple and ring.
- Once feeding begins, use within 1 hour or discard.

#### Ready to Feed

Your baby's health depends on carefully following these directions. Ask your baby's doctor if you need to boil (sterilize) nipples and rings before use. Ready to feed formula if mixed with water may not provide proper nutrition, and repeated use of such feedings could cause illness.

- 1 Do not add water.
- 1 Shake very well.

### 2 or 8 fl oz bottle

- 1 Do not use if breakaway ring is missing or broken.
- Twist off cap.
- 1 Use clean hands to twist on nipple and ring. Use sterile nipple and ring if directed.
- For use in a different feeding container (cup or bottle):
  - 1 Ask your baby's doctor if you need to boil (sterilize) the empty bottles before adding formula.
  - 1 Pour desired amount into clean feeding container. Recap bottle, refrigerate and use within 48 hours.
- 1 Once feeding begins, use within 1 hour or discard.

#### 32 fl oz bottle

- 1 Do not use if band around cap or inner foil seal is damaged.
- 1 Shake very well before each use.
- Remove protective band; twist off and clean cap.
- 1 Invert cap; press down to pierce foil, then turn cap a half turn.
- 1 Remove foil.
- 1 Pour formula into bottle; attach clean nipple.
- Once feeding begins, use within 1 hour or discard.
- 1 Prepare and feed all formula in the container within 3 days after opening.

#### Storage Instructions:

#### Powder

#### Cans and SimplePac

- Once mixed, store bottles in refrigerator and feed to baby within 24 hours.
- Store unopened conainers at room temperature; avoid extreme temperatures.
- Use opened can contents within 1 month.
- 1 Do not reuse container.

Abbott

### Milk-based infant formula with iron

#### Powder Singles

- Once mixed, feed immediately or cover and refrigerate; use within 24 hours.
- Store unopened packets at room temperature; avoid extreme temperatures.

#### Ready to Feed

#### 2 & 8 fl oz Bottle

- 1 Store unopened at room temperature; avoid extreme temperatures.
- 1 Do not reuse bottle.
- 1 Protect from light.
- Store in carton until ready for use.

#### 32 fl oz Bottle

- 1 Once opened, store quart bottle in refrigerator.
- 1 Store prepared bottles in refrigerator and feed to baby within 48 hours.
- 1 Store unopened containers at room temperature; avoid extreme temperatures.
- 1 Do not reuse container.

