

Product Information: Similac® For Spit-Up NON-GMO

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Similac® For Spit-Up NON-GMO

Infant Formula with Iron

- 1 A 19 Cal/fl oz nutritionally complete, non-GMO, milk-based infant formula with added rice starch to help reduce frequent spit up.
- 1 Suitable for lactose sensitivity.*
- 1 Non-GMO. Ingredients not genetically engineered.
- 1 A non-GMO[†] option for complete nutrition for baby's first year.
- 1 OptiGRO™ is our exclusive blend of DHA, lutein and vitamin E: these important ingredients are found in breast milk.
 - DHA for brain and eye development.
 - Lutein to support eye health.
 - Vitamin E, an important nutrient to support developing cells.
- 1 Our exclusive formula has:
 - Lutein and DHA for baby's developing eyes and brain.
 - Calcium for strong bones - no palm olein oil.
 - Nucleotides to help support the immune system.
 - Prebiotics to help promote digestive health.
 - Carotenoids like those naturally found in breast milk.
- 1 Easy to digest milk-based formula with added rice starch.
- 1 Clinically shown to reduce frequent spit up in healthy babies by 54%.^{1,‡}
- 1 Clinically shown to support normal infant growth.¹
- 1 Kosher.
- 1 Halal.
- 1 Gluten-free.



Safety Precautions

- 1 **Never use a microwave oven to warm formula.** Serious burns can result.
- 1 Not for use by infants or children with galactosemia.
- 1 Powdered infant formulas are not sterile and should not be fed to premature infants or infants who might have immune problems unless directed and supervised by your baby's doctor.

* Typical value 0.1 g lactose/100 kcal compared to Similac Advance (11 g lactose/100 kcal).

[†] Ingredients not genetically engineered.

[‡] Among healthy 2-month-old infants compared to a standard formula.

¹ Lasekan JB, et al. *J Am Coll Nutr.* 2014;33(2):136-146.

Ingredients

Powder Unflavored:

Corn Syrup, Modified Rice Starch, Milk Protein Isolate, High Oleic Safflower Oil, Sugar, Soy Oil, Coconut Oil, Galactooligosaccharides. Less than 2% of: C. Cohnii Oil, M. Alpina Oil, Beta-Carotene, Lutein, Lycopene, Calcium Phosphate, Potassium Citrate, Potassium Chloride, Potassium Phosphate, Calcium Carbonate, Magnesium Chloride, Ascorbic Acid, Choline Chloride, L-Cystine Dihydrochloride, Salt, Ascorbyl Palmitate, Ferrous Sulfate, Taurine, m-Inositol, Choline Bitartrate, Zinc Sulfate, L-Carnitine, Mixed Tocopherols, Niacinamide, d-Alpha-Tocopheryl Acetate, Calcium Pantothenate, Vitamin A Palmitate, Copper Sulfate, Thiamine Hydrochloride, Riboflavin, Pyridoxine Hydrochloride, Folic Acid, Potassium Iodide, Phylloquinone, Biotin, Sodium Selenate, Vitamin D3, Vitamin B12, Potassium Hydroxide, and Nucleotides (Adenosine 5'-Monophosphate, Cytidine 5'-Monophosphate, Disodium Guanosine 5'-Monophosphate, Disodium Uridine 5'-Monophosphate).

Allergens: Contains milk ingredients.

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Similac® For Spit-Up NON-GMO

Infant Formula with Iron

Availability

List Number	Item
53729	Similac for Spit Up NON-GMO Powder / 22.5 oz (638 g) SimplePac / 6 ct
56728	Similac for Spit Up Ready to Feed / 1 QT (946 mL) Bottle / 6 ct
62139	Similac for Spit Up NON-GMO Ready to Feed / 2 fl oz (59 mL) bottle / 48 ct

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Similac® For Spit-Up NON-GMO

Infant Formula with Iron

Nutrition Information - Powder Unflavored

	100 Cal (5.3 fl oz, prepared as directed)		1000 mL (1.1 QT, prepared as directed)	
	Value	%RDI / %DV	Value	%RDI / %DV
Calories	100		643	
Volume, mL	155.5		1000	
Protein, g	2.14		13.76	
Fat, g	5.40		34.73	
Carbohydrate, g	11		70.74	
Water, g	140		902	
Linoleic Acid, mg	1000		6431	
Vitamin A, IU	300		1929	
Vitamin D, IU	60		386	
Vitamin E, IU	1.5		9.6	
Vitamin K, mcg	8		51.4	
Thiamin (Vitamin B1), mcg	100		643	
Riboflavin (Vitamin B2), mcg	160		1029	
Vitamin B6, mcg	63		405	
Vitamin B12, mcg	0.26		1.67	
Niacin, mcg	1100		7074	
Folic Acid (Folacin), mcg	16		102.9	
Pantothenic Acid, mcg	470		3023	
Biotin, mcg	4.6		29.6	
Vitamin C (Ascorbic Acid), mg	9		57.9	
Choline, mg	24		154.3	
Inositol, mg	4.5		28.9	
Calcium, mg	88		566	
Calcium, mEq	4.4		28.3	
Phosphorus, mg	59		379	
Magnesium, mg	6		38.6	
Iron, mg	1.9		12.22	
Zinc, mg	0.79		5.08	
Manganese, mcg	5		32.2	
Copper, mcg	95		611	
Iodine, mcg	9		58	
Selenium, mcg	2		13	
Sodium, mg	32		206	
Sodium, mEq	1.4		9	
Potassium, mg	110		707	
Potassium, mEq	2.8		18	
Chloride, mg	68		437	
Chloride, mEq	1.9		12.2	

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Similac® For Spit-Up NON-GMO

Infant Formula with Iron

Preparation

Powder

Your baby's health depends on carefully following these directions. Proper hygiene, handling and storage are important when preparing infant formula. Failure to follow these directions could result in severe harm. Ask your baby's doctor if you need to use cooled, boiled water for mixing and if you need to boil (sterilize) bottles, nipples and rings before use.

- 1 Wash your hands, surfaces and utensils.
- 1 Pour water into clean bottle (see mixing guide).
- 1 Add 1 unpacked level scoop (8.3 g) to each 2 fl oz of water.
- 1 Return dry scoop to container.
- 1 Cap bottle; shake well; wait two minutes; attach nipple; gently swirl before feeding.
- 1 Once oral feeding begins, use **within 1 hour or discard**.

Mixing Guide

Measure water	+	Add scoop(s) of <i>unpacked level powder</i> using enclosed scoop	=	Finished bottle (approx)
2 fl oz		1 scoop (8.3 g)		2 fl oz
4 fl oz		2 scoops		4 fl oz
6 fl oz		3 scoops		6 fl oz
8 fl oz		4 scoops		8 fl oz

Ready to Feed

Your baby's health depends on carefully following these directions. Failure to follow these directions could result in severe harm. Ask your baby's doctor if you need to boil (sterilize) bottles, nipples and rings before use.

2-fl-oz Bottle

- 1 Do not add water.
- 1 Shake very well.
- 1 **Do not use if breakaway ring is missing or broken.**
- 1 Twist off cap.
- 1 Twist on nipple and ring.
 - 1 Use sterile nipple and ring if directed.
- 1 Once oral feeding begins, use **within 1 hour or discard**.

1-QT Bottle

- 1 Do not add water.
- 1 Shake very well before each use.
- 1 **Do not use if band around cap or inner foil seal is damaged.**
- 1 Remove protective band; twist off and clean cap.
- 1 Invert cap; press down to piece foil, then turn cap a half turn.
- 1 Remove foil.
- 1 Pour formula into bottle; attach nipple.
- 1 Once oral feeding begins, use **within 1 hour or discard**.
- 1 Prepare and feed all formula from the container **within 3 days after opening**.

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Storage & Handling

Powder

- 1 Once mixed, store bottles in refrigerator and **feed to baby within 24 hours.**
- 1 Store unopened or opened container at room temperature; avoid extreme temperatures.
- 1 Use opened can contents **within 1 month.**
- 1 Do not reuse container.

Ready to Feed

2-fl-oz Bottle

- 1 Avoid prolonged exposure of bottles to light.
- 1 Store unopened containers at room temperature; avoid extreme temperatures.

1-QT Bottle

- 1 Once opened, store quart bottle in refrigerator.
- 1 Store prepared bottles in refrigerator and **feed to baby within 48 hours.**
- 1 Store unopened containers at room temperature; avoid extreme temperatures.
- 1 Prepare and feed all formula from this container **within 3 days after opening.**
- 1 Do not reuse container.