

Product Information: Similac® For Supplementation

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Similac® For Supplementation

Infant formula with iron

- 1 A 19 Cal/fl oz, nutritionally complete, milk-based, iron-fortified, non-GMO[†] formula for breastfeeding moms who choose to introduce formula. For a gentle introduction to formula.
- 1 No artificial growth hormones.[†]
- 1 OptiGRO™ is our exclusive blend of DHA, lutein and vitamin E: these important ingredients are found in breast milk.
 - DHA for brain and eye development.
 - Lutein to support eye health.
 - Vitamin E, an important nutrient to support developing cells.
- 1 Our exclusive formula has:
 - Calcium for strong bones - no palm olein oil.
 - Nucleotides to help support the immune system.
 - Prebiotics to help promote digestive health.
 - Carotenoids like those naturally found in breast milk.
- 1 Gluten-free.
- 1 Kosher.
- 1 Halal.



Safety Precautions

- 1 **Never use a microwave oven to warm formula.** Serious burns can result.
- 1 Powdered infant formulas are not sterile and should not be fed to premature infants or infants who might have immune problems unless directed and supervised by your baby's doctor.
- 1 Not for use by infants or children with galactosemia.

* *Ingredients not genetically engineered.*

† *No significant difference has been shown between milk derived from rbST-treated and non-rbST-treated cows.*

Ingredients

Powder Unflavored:

Nonfat Milk, Lactose, Whey Protein Concentrate, High Oleic Safflower Oil, Soy Oil, Coconut Oil, Galactooligosaccharides. Less than 2% of: C. Cohnii Oil, M. Alpina Oil, Beta-Carotene, Lutein, Lycopene, Potassium Citrate, Calcium Carbonate, Soy Lecithin, Ascorbic Acid, Potassium Chloride, Magnesium Chloride, Ferrous Sulfate, Choline Bitartrate, Choline Chloride, Ascorbyl Palmitate, Salt, Taurine, m-Inositol, Zinc Sulfate, Mixed Tocopherols, d-Alpha-Tocopheryl Acetate, Niacinamide, Calcium Pantothenate, L-Carnitine, Vitamin A Palmitate, Cupric Sulfate, Thiamine Chloride Hydrochloride, Riboflavin, Pyridoxine Hydrochloride, Folic Acid, Manganese Sulfate, Phylloquinone, Biotin, Sodium Selenate, Vitamin D3, Cyanocobalamin, Calcium Phosphate, Potassium Phosphate, Potassium Hydroxide, and Nucleotides (Adenosine 5'-Monophosphate, Cytidine 5'-Monophosphate, Disodium Guanosine 5'-Monophosphate, Disodium Uridine 5'-Monophosphate).

Allergens: Contains milk and soy ingredients.

Availability

List Number	Item
62990	Similac For Supplementation Ready to Feed / 2 fl oz (59 mL) Bottle / 6 x 8 ct
63013	Similac For Supplementation Powder / 23.2 oz (658 g) SimplePac / 4 ct
63092	Similac For Supplementation Ready to Feed / 2 fl oz (59 mL) Bottle / 48 ct
67199	Similac For Supplementation Ready to Feed / 2 fl oz (59 mL) Bottle / 8 x 6 ct
67440	Similac For Supplementation Ready to Feed / 2 fl oz (59 mL) Bottle / 12 x 4 ct

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Abbott Nutrition Abbott Laboratories
Columbus, OH 43219-3034
1-800-227-5767



Similac® For Supplementation

Infant formula with iron

Nutrition Information - Powder Unflavored

	100 Cal (5.3 fl oz, prepared as directed)		1000 mL (1.1 QT, prepared as directed)	
	Value	%RDI / %DV	Value	%RDI / %DV
Calories	100		643	
Volume, mL	156		1000	
Protein, g	2.07		13.31	
Fat, g	5.60		36.08	
Carbohydrate, g	10.8		69.58	
Water, g	141		907	
Linoleic Acid, mg	1000		6431	
Vitamin A, IU	300		1929	
Vitamin D, IU	75		482	
Vitamin E, IU	1.5		9.6	
Vitamin K, mcg	8		51	
Thiamin (Vitamin B1), mcg	100		643	
Riboflavin (Vitamin B2), mcg	160		1029	
Vitamin B6, mcg	63		405	
Vitamin B12, mcg	0.26		1.67	
Niacin, mcg	1100		7074	
Folic Acid (Folacin), mcg	16		103	
Pantothenic Acid, mcg	470		3023	
Biotin, mcg	4.6		29.6	
Vitamin C (Ascorbic Acid), mg	9		58	
Choline, mg	24		154	
Inositol, mg	4.9		31.5	
Calcium, mg	82		527	
Calcium, mEq	4.1		26.4	
Phosphorus, mg	44		283	
Magnesium, mg	6		39	
Iron, mg	1.9		12.2	
Zinc, mg	0.79		5.08	
Manganese, mcg	5		32	
Copper, mcg	95		611	
Iodine, mcg	6		39	
Selenium, mcg	2		13	
Sodium, mg	25		161	
Sodium, mEq	1.1		7.1	
Potassium, mg	110		707	
Potassium, mEq	2.8		18	
Chloride, mg	68		437	
Chloride, mEq	1.9		12.2	

For more information, contact your Abbott Nutrition Representative or visit www.abbottnutrition.com

Similac® For Supplementation

Infant formula with iron

Preparation

Instructions for Use:

Your baby's health depends on carefully following these directions. Proper hygiene, handling and storage are important when preparing infant formula. Failure to follow these directions could result in severe harm. Ask your baby's doctor if you need to use cooled, boiled water for mixing and if you need to boil (sterilize) bottle, nipple and ring before use.

Powder

- 1 Wash your hands, surfaces and utensils.
- 1 Pour water into clean bottle (see Powder Mixing guide).
- 1 Add 1 unpacked level scoop (8.3 g) to each 2 fl oz of water.
- 1 Return dry scoop to holder in lid.
- 1 Cap bottle; shake well; attach nipple.
- 1 Once feeding begins, **use within 1 hour or discard.**

Powder Mixing Guide

Measure water	+	Add scoop(s) of <i>unpacked level powder</i> using enclosed scoop	=	Finished bottle (approx)
2 fl oz		1 scoop (8.3 g)		2 fl oz
4 fl oz		2 scoops		4 fl oz
6 fl oz		3 scoops		6 fl oz
8 fl oz		4 scoops		8 fl oz

Ready to Feed

Ready to feed formula if mixed with water may not provide proper nutrition, and repeated use of such feedings could cause illness.

- 1 **Do not use if breakaway ring is missing or broken.**
- 1 **Do not add water.**
- 1 **Shake very well.**
- 1 Twist off cap.
- 1 Twist on nipple and ring.
 - 1 Since this formula is commercially sterile, use clean hands to attach any clean standard nipple and ring.
 - 1 Use sterile nipple and ring if directed.
- 1 No need to warm before feeding.
- 1 **Once oral feeding begins, use within 1 hour or discard.**

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Abbott Nutrition Abbott Laboratories
Columbus, OH 43219-3034
1-800-227-5767



Storage & Handling

Instructions for Use:

Powder

- 1 Once mixed, store bottles in refrigerator and **feed to baby within 24 hours.**
- 1 Store unopened or opened container at room temperature; avoid extreme temperatures.
- 1 **Use opened can contents within 1 month.**
- 1 Do not reuse container.

Ready to Feed

- 1 Store unopened at room temperature; avoid extreme temperatures.
- 1 Do not reuse bottle.
- 1 **Protect from light.**
- 1 **Store in carton until ready for use.**