Infant Formula with Iron

Product Information: Similac Pro-Advance®



Infant Formula with Iron

- 1 A 20 Cal/fl oz, nutritionally complete, non-GMO,* milk-based, iron-fortified standard infant formula designed to support both immune and cognitive development with key ingredients found in breast milk. For use as a supplement or alternative to breastfeeding.
- SIMILAC PRO-ADVANCE with 2'-FL Human Milk Oligosaccharide[†] helps strengthen the immune system to be more like the breastfed infant than ever before. 1,2,3
- Nucleotides to help support the developing immune system.^{2,3}
- OptiGRO® is our exclusive blend of DHA, lutein and vitamin E: these important ingredients are found in breast milk.
 - O DHA for brain and eye development.
 - Lutein to support eye health.
 - O Vitamin E; an important nutrient found in breast milk to support developing cells.
- 1 Our exclusive formula has:
 - O Calcium for strong bones no palm olein oil.
 - O Prebiotics to help promote digestive health.
 - O Carotenoids like those naturally found in breast milk.
- Clinically shown to support normal growth and tolerance in newborns. 4,5
- 1 Clinically shown to support greater calcium absorption[‡] for strong bones.^{6,7}
- 1 Non-GMO.
- I Gluten-free.
- 1 Halal.
- 1 Kosher made with dairy ingredients.

Safety Precautions

- Not for infants or children with galactosemia.
- Never use a microwave oven to warm formula. Serious burns can result.
- 1 Powdered infant formulas are not sterile and should not be fed to premature infants or infants who might have immune problems unless directed and supervised by your baby's doctor.

Ingredients



^{*} Ingredients not genetically engineered.

^{*} Ingredients not genetically engineered.

[†] Not from human milk.

[‡] Similac® with Iron vs Enfamil® with Iron. Enfamil is not a trademark of Abbott Laboratories.

¹ Goehring KC, et al. J Nutr. 2016;146(12):2559-2566.

² Schaller JP, et al. Pediatr Res. 2004;56(6):883-890.

³ Buck RH, et al. Pediatr Res. 2004;56(6):891-900.

⁴ Marriage B, et al. JPGN 2015;61:649-653.

⁵ Kajzer J, et al. FASEB J. 2016;30(Supp 1):671-674.

⁶ Nelson SE, et al. J Am Coll Nutr. 1998;17(4):327-332.

⁷ Koo W, et al. Pediatrics. 2003;111:1017-1023.

Infant Formula with Iron

Powder Unflavored:

Nonfat Milk, Lactose, High Oleic Safflower Oil, Whey Protein Concentrate, Soy Oil, Coconut Oil. Less than 2% of: Schizochytrium Sp. Oil*, M. Alpina Oil[†], 2'-Fucosyllactose[‡], Short-chain Fructooligosaccharides, Beta-Carotene, Lutein, Potassium Citrate, Calcium Carbonate, Ascorbic Acid, Soy Lecithin, Potassium Chloride, Magnesium Chloride, Ferrous Sulfate, Choline Bitartrate, Choline Chloride, Ascorbyl Palmitate, Salt, Taurine, Inositol, Zinc Sulfate, Mixed Tocopherols, d-Alpha-Tocopheryl Acetate, Niacinamide, Calcium Pantothenate, L-Carnitine, Vitamin A Palmitate, Copper Sulfate, Thiamine Hydrochloride, Riboflavin, Pyridoxine Hydrochloride, Folic Acid, Manganese Sulfate, Phylloquinone, Biotin, Sodium Selenate, Vitamin D3, Vitamin B12, Calcium Phosphate, Potassium Phosphate, Potassium Iodide, Potassium Hydroxide, and Nucleotides (Adenosine 5'-Monophosphate, Cytidine 5'-Monophosphate, Disodium Guanosine 5'-Monophosphate, Disodium Uridine 5'-Monophosphate).

Allergens: Contains milk and soy ingredients.

Availability

List Number	Item		
66439	Similac Pro-Advance Powder / 30.8 oz (873 g) Can / 4 ct		
66655	Similac Pro-Advance Powder / 34 oz (964 g) SimplePac / 6 ct		
68088	Similac Pro-Advance Powder / 20.6 oz (584 g) Simplepac/ 4 ct		

^{*} Source of DHA.

[†] Source of ARA.

[‡] 2'-FL, is derived from milk, but does not contain milk protein.

Infant Formula with Iron

Nutrition Information - Powder Unflavored

Nutrition Information - Powder Unflavored						
	100 Cal (5 fl oz, p	100 Cal (5 fl oz, prepared as directed)		1000 mL (1.1 QT, prepared as directed)		
	Value	%RDI / %DV	Value	%RDI / %DV		
Calories	100		676			
Volume, mL	148		1000			
Protein, g	2.07		14			
Fat, g	5.6		37.87			
Carbohydrate, g	10.5		71.01			
Water, g	134		900			
Linoleic Acid, mg	1000		6763			
Vitamin A, IU	300		2029			
Vitamin D, IU	60		406			
Vitamin E, IU	1.5		10.1			
Vitamin K, mcg	8		54.1			
Thiamin (Vitamin B1), mcg	100		676			
Riboflavin (Vitamin B2), mcg	150		1014			
Vitamin B6, mcg	60		406			
Vitamin B12, mcg	0.25		1.69			
Niacin, mcg	1050		7101			
Folic Acid (Folacin), mcg	15		101.4			
Pantothenic Acid, mcg	450		3043			
Biotin, mcg	4.4		29.8			
Vitamin C (Ascorbic Acid), mg	9		60.9			
Choline, mg	24		162.3			
Inositol, mg	24		162.3			
Calcium, mg	78		527			
Calcium, mEq	3.9		26.4			
Phosphorus, mg	42		284			
Magnesium, mg	6		40.6			
Iron, mg	1.8		12.17			
Zinc, mg	0.75		5.07			
Manganese, mcg	5		33.8			
Copper, mcg	90		609			
lodine, mcg	15		101			
Selenium, mcg	2		14			
Sodium, mg	24		162			
Sodium, mEq	1		6.8			
Potassium, mg	105		710			
Potassium, mEq	2.7		18.3			
Chloride, mg	65		440			
Chloride, mEq	1.8		12.2			



Infant Formula with Iron

Preparation

Instructions for Use:

Your baby's health depends on carefully following these directions. Proper hygiene, handling and storage are important when preparing infant formula. Failure to follow these directions could result in severe harm. Ask your baby's doctor if you need to use cooled, boiled water for mixing and if you need to boil (sterilize) bottles, nipples and rings before use.

- 1 Wash your hands, surfaces and utensils.
- Pour water into clean bottle.
- 1 Add 1 unpacked level scoop (8.6 g) to each 2 fl oz of water.
- 1 Return dry scoop to container.
- 1 Cap bottle; shake well; attach nipple.
- 1 Once feeding begins, use within 1 hour or discard.

Storage Instructions:

- 1 Once mixed, store bottles in refrigerator and feed to baby within 24 hours.
- 1 Store unopened containers at room temperature; avoid extreme temperatures.
- Use opened can contents within 1 month.
- 1 Do not reuse container.

