

## Product Information: Similac Pro-Sensitive®

For more information, contact your Abbott Nutrition Representative or visit  
[www.abbottnutrition.com](http://www.abbottnutrition.com)

# Similac Pro-Sensitive®

## Infant formula with iron



- 1 A 20 Cal/fl oz, nutritionally complete, non-GMO,<sup>\*</sup> iron-fortified infant formula designed for use as a milk-based alternative to standard milk-based formulas for sensitive tummies.
- 1 Gentle nutrition designed to ease common tummy troubles like fussiness,<sup>†</sup> gas,<sup>†</sup> or mild spit-up
- 1 Designed to support both immune and cognitive development with key ingredients found in breast milk.
- 1 Suitable for infants with lactose sensitivity.<sup>‡</sup>
- 1 Similac Pro-Sensitive 20 Cal/fl oz with 2'-FL Human Milk Oligosaccharide,<sup>§</sup> helps strengthen the immune system to be more like the breastfed infant than ever before.<sup>1,2,3</sup>
- 1 Nucleotides to help support the developing immune system.<sup>2,3</sup>
- 1 OptiGRO® is our exclusive blend of DHA, lutein and vitamin E: these important ingredients are found in breast milk.
  - DHA for brain and eye development.
  - Lutein to support eye health.
  - Vitamin E, an important nutrient to support developing cells.
- 1 Our exclusive formula has:
  - Calcium for strong bones - no palm olein oil.
  - Prebiotics to help promote digestive health.
  - Carotenoids like those naturally found in breast milk.
- 1 Promotes growth, tolerance, and soft stools.
- 1 Low osmolality (200 mOsm/kg water).
- 1 Non-GMO.<sup>\*</sup>
- 1 No artificial growth hormones.<sup>¶</sup>
- 1 Gluten-free.
- 1 Halal.
- 1 Kosher made with dairy ingredients.

## Safety Precautions

- 1 **Never use a microwave oven to warm formula.** Serious burns can result.
- 1 Not for infants or children with galactosemia.

<sup>\*</sup> *Ingredients not genetically engineered.*

<sup>\*</sup> *Ingredients not genetically engineered.*

<sup>†</sup> *Due to lactose sensitivity.*

<sup>‡</sup> *Typical value 0.1 g lactose/100 kcal compared to Similac® Advance® (11 g lactose/100 kcal).*

<sup>§</sup> *Not from human milk.*

<sup>¶</sup> *No significant difference has been shown between milk derived from rbST-treated and non-rbST-treated cows.*

<sup>1</sup> *Goehring KC, et al. J Nutr. 2016;146(12):2559-2566.*

<sup>2</sup> *Schaller JP, et al. Pediatr Res. 2004;56(6):883-890.*

<sup>3</sup> *Buck RH, et al. Pediatr Res. 2004;56(6):891-900.*

## Ingredients

For more information, contact your Abbott Nutrition Representative or visit  
[www.abbottnutrition.com](http://www.abbottnutrition.com)

# Similac Pro-Sensitive®

Infant formula with iron

## Powder Unflavored:

Corn Syrup, Milk Protein Isolate, High Oleic Safflower Oil, Sugar, Soy Oil, Coconut Oil. Less than 2% of: C. Cohnii Oil, M. Alpina Oil, 2'-Fucosyllactose, Short-chain Fructooligosaccharides, Beta-Carotene, Lutein, Potassium Chloride, Sodium Citrate, Calcium Phosphate, Potassium Citrate, Soy Lecithin, Inositol, Magnesium Phosphate, Ascorbic Acid, Choline Chloride, Ascorbyl Palmitate, Ferrous Sulfate, Choline Bitartrate, Taurine, Calcium Carbonate, Zinc Sulfate, L-Carnitine, Mixed Tocopherols, Niacinamide, d-Alpha-Tocopheryl Acetate, Calcium Pantothenate, Magnesium Chloride, Vitamin A Palmitate, Copper Sulfate, Thiamine Hydrochloride, Riboflavin, Pyridoxine Hydrochloride, Folic Acid, Potassium Iodide, Manganese Sulfate, Phylloquinone, Biotin, Sodium Selenate, Vitamin D3, Vitamin B12, Potassium Hydroxide, and Nucleotides (Adenosine 5'-Monophosphate, Cytidine 5'-Monophosphate, Disodium Guanosine 5'-Monophosphate, Disodium Uridine 5'-Monophosphate).

**Allergens:** Contains milk and soy ingredients.

## Availability

List Number	Item
64253	Similac Pro-Sensitive Ready to Feed / 32 fl oz (946 mL) Bottle / 6 ct
66441	Similac Pro-Sensitive Powder / 29.8 oz (845 g) Can / 4 ct
66657	Similac Pro-Sensitive Powder / 34 oz (964 g) SimplePac / 6 ct
67325	Similac Pro-Sensitive Ready to Feed / 2 fl oz (59 mL) Bottle / 12 x 4 ct
67517	Similac Pro-Sensitive Ready to Feed / 8 fl oz (237 mL) Bottle / 4 x 6 ct
67930	Similac Pro-Sensitive Powder / 29.8 oz (845 g) Can / 6 ct

For more information, contact your Abbott Nutrition Representative or visit  
[www.abbottnutrition.com](http://www.abbottnutrition.com)

### Nutrition Information - Powder Unflavored

	100 Cal (5 fl oz, prepared as directed)		1000 mL (1.1 QT, prepared as directed)	
	Value	%RDI / %DV	Value	%RDI / %DV
Calories	100		676	
Volume, mL	148		1000	
Protein, g	2.1		14.19	
Fat, g	5.4		36.49	
Carbohydrate, g	10.9		73.65	
Water, g	133		899	
Linoleic Acid, mg	1000		6757	
Vitamin A, IU	300		2027	
Vitamin D, IU	60		405	
Vitamin E, IU	1.5		10.1	
Vitamin K, mcg	8		54.1	
Thiamin (Vitamin B1), mcg	100		676	
Riboflavin (Vitamin B2), mcg	150		1014	
Vitamin B6, mcg	60		405	
Vitamin B12, mcg	0.25		1.69	
Niacin, mcg	1050		7095	
Folic Acid (Folacin), mcg	15		101.4	
Pantothenic Acid, mcg	450		3041	
Biotin, mcg	4.4		29.7	
Vitamin C (Ascorbic Acid), mg	9		60.8	
Choline, mg	24		162.2	
Inositol, mg	24		162.2	
Calcium, mg	84		568	
Calcium, mEq	4.2		28.4	
Phosphorus, mg	56		378	
Magnesium, mg	6		40.5	
Iron, mg	1.8		12.16	
Zinc, mg	0.75		5.07	
Manganese, mcg	5		33.8	
Copper, mcg	90		608	
Iodine, mcg	15		101	
Selenium, mcg	2		14	
Sodium, mg	30		203	
Sodium, mEq	1.3		8.8	
Potassium, mg	107		723	
Potassium, mEq	2.7		18.2	
Chloride, mg	65		439	
Chloride, mEq	1.8		12.2	

For more information, contact your Abbott Nutrition Representative or visit  
[www.abbottnutrition.com](http://www.abbottnutrition.com)

### Preparation

#### Instructions for Use:

##### Powder

Your baby's health depends on carefully following these directions. Proper hygiene, handling and storage are important when preparing infant formula. Failure to follow these directions could result in severe harm. Ask your baby's doctor if you need to use cooled, boiled water for mixing and if you need to boil (sterilize) bottles, nipples and rings before use.

- 1 Wash your hands, surfaces and utensils.
- 1 Pour water into clean bottle.
- 1 Add 1 unpacked level scoop (8.8 g) to each 2 fl oz of water.
- 1 Return dry scoop to container.
- 1 Cap bottle; shake well; attach nipple.
- 1 Once feeding begins, use within 1 hour or discard.

##### Ready to Feed:

Your baby's health depends on carefully following these directions. Failure to follow these directions could result in severe harm. Ask your baby's doctor if you need to boil (sterilize) bottle, nipple and ring before use. Ready to feed formula if mixed with water may not provide proper nutrition, and repeated use of such feedings could cause illness.

- 1 **Use as directed by a doctor.**
- 1 **Do not add water.**

##### 2 or 8 fl oz bottle

- 1 **Do not use if breakaway ring is missing or broken.**
- 1 Shake very well.
- 1 Twist off cap.
- 1 Twist on nipple and ring.
- 1 Use clean hands to attach nipple and ring. Use sterile nipple and ring if directed.
- 1 For use in a different feeding container (cup or bottle):
  - 1 Pour desired amount into clean feeding container.
  - 1 Recap bottle, refrigerate and use within 48 hours.
- 1 Once oral feeding begins, use within 1 hour or discard.

##### 1 QT bottle

- 1 **Do not use if band around cap or inner foil seal is damaged.**
- 1 Shake very well before each use.
- 1 Remove protective band, twist off and clean cap.
- 1 Invert cap; press down to pierce foil, then turn cap a half turn.
- 1 Remove foil.
- 1 Pour formula into bottle; attach nipple.
- 1 Once feeding begins, use within 1 hour or discard.
- 1 Prepare and feed all formula in the container within 3 days after opening.

## Storage & Handling

### Instructions for Use:

#### Powder

- 1 Once mixed, store bottles in refrigerator and **feed to baby within 24 hours.**
- 1 Store unopened or opened container at room temperature; avoid extreme temperatures.
- 1 **Use opened container contents within 1 month.**
- 1 Do not reuse container.

#### Ready to Feed

##### 2 or 8 fl oz bottles

- 1 Store unopened at room temperature; avoid extreme temperatures.
- 1 **Protect from light.**
- 1 **Store 2 fl oz bottles in carton until ready for use.**
- 1 Do not reuse container.

##### 1 QT bottle

- 1 Once opened, store quart bottle in refrigerator.
- 1 Store prepared bottles in refrigerator and **feed to baby within 48 hours.**
- 1 Store unopened containers at room temperature; avoid extreme temperatures.
- 1 Do not reuse container.