

Product Information: Similac Sensitive® Liquid

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Similac Sensitive® Liquid

Infant formula with iron

- 1 SIMILAC SENSITIVE is a 20 Cal/fl oz, nutritionally complete infant formula that is an alternative to standard milk-based formulas. For fussiness,* gas,* or mild spit-up. Suitable for infants with lactose sensitivity.†
- 1 Easy-to-digest gentle carbohydrate for fussiness and gas due to lactose sensitivity.
- 1 OptiGRO® is our exclusive blend of DHA, lutein and vitamin E: these important ingredients are found in breast milk.
 - DHA for brain and eye development.
 - Lutein to support eye health.
 - Vitamin E to support developing cells.
- 1 Our exclusive formula has:
 - Calcium for strong bones - no palm olein oil.
 - Nucleotides to help support the immune system.
 - Prebiotics to help promote digestive health.
 - Carotenoids like those naturally found in breast milk.
- 1 Clinically shown to support normal growth in infants.¹
- 1 A unique blend of carbohydrates using two different absorptive pathways to help maximize absorption and minimize malabsorption risks.
- 1 Low osmolality (200 mOsm/kg water).
- 1 Kosher made with dairy ingredients.
- 1 Halal.



Safety Precautions

- 1 **Never use a microwave oven to warm formula.** Serious burns can result.
- 1 Not for infants or children with galactosemia.

* Due to lactose sensitivity.

† Typical value 0.1 g lactose/100 kcal compared to Similac® Advance® (11 g lactose/100 kcal).

¹ Lasekan JB, et al. Clin Pediatr. 2011;50(4):330-337.

Ingredients

Liquid Unflavored:

Water, Maltodextrin, Milk Protein Isolate, High Oleic Safflower Oil, Sugar, Soy Oil, Coconut Oil. Less than 0.5% of: Schizochytrium Sp. Oil*, M. Alpina Oil†, Short-chain Fructooligosaccharides‡, Beta-Carotene, Lutein, Calcium Phosphate, Potassium Citrate, Carrageenan, Potassium Chloride, Ascorbic Acid, Monoglycerides, Soy Lecithin, Magnesium Chloride, Calcium Carbonate, Inositol, Choline Chloride, Magnesium Phosphate, Sodium Citrate, Ferrous Sulfate, Taurine, Choline Bitartrate, Zinc Sulfate, L-Carnitine, Niacinamide, d-Alpha-Tocopheryl Acetate, Calcium Pantothenate, Riboflavin, Copper Sulfate, Vitamin A Palmitate, Thiamine Hydrochloride, Pyridoxine Hydrochloride, Folic Acid, Potassium Iodide, Manganese Sulfate, Phylloquinone, Biotin, Sodium Selenate, Vitamin D3, Vitamin B12, Potassium Hydroxide, and Nucleotides (Adenosine 5'-Monophosphate, Cytidine 5'-Monophosphate, Disodium Guanosine 5'-Monophosphate, Disodium Uridine 5'-Monophosphate).

* Source of DHA.

† Source of ARA.

‡ Prebiotic.

Allergens: Contains milk and soy ingredients.

Availability

List Number	Item
57533	Similac Sensitive Ready to Feed / 32 fl oz (946 mL) Bottle / 6 ct
57535	Similac Sensitive Concentrated Liquid / 13 fl oz (384 mL) Can / 6 ct

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Abbott Nutrition Abbott Laboratories
Columbus, OH 43219-3034
1-800-227-5767

2 of 4



Similac Sensitive® Liquid

Infant formula with iron

Nutrition Information - Liquid Unflavored

	100 Cal (5 fl oz)		1000 mL (1.1 QT)	
	Value	%RDI / %DV	Value	%RDI / %DV
Calories	100		676	
Volume, mL	148		1000	
Protein, g	2.1		14.19	
Fat, g	5.4		36.49	
Carbohydrate, g	10.9		73.65	
Water, g	133		899	
Linoleic Acid, mg	1000		6757	
Potential Renal Solute Load, mOsm	19.7		132.8	
Vitamin A, IU	300		2027	
Vitamin D, IU	60		405	
Vitamin E, IU	1.5		10.1	
Vitamin K, mcg	8		54.1	
Thiamin (Vitamin B1), mcg	100		676	
Riboflavin (Vitamin B2), mcg	150		1014	
Vitamin B6, mcg	60		405	
Vitamin B12, mcg	0.25		1.69	
Niacin, mcg	1050		7095	
Folic Acid (Folacin), mcg	15		101.4	
Pantothenic Acid, mcg	450		3041	
Biotin, mcg	4.4		29.7	
Vitamin C (Ascorbic Acid), mg	9		60.8	
Choline, mg	24		162.2	
Inositol, mg	24		162.2	
Calcium, mg	84		568	
Calcium, mEq	4.2		28.4	
Phosphorus, mg	56		378	
Magnesium, mg	6		40.5	
Iron, mg	1.8		12.16	
Zinc, mg	0.75		5.07	
Manganese, mcg	5		33.8	
Copper, mcg	90		608	
Iodine, mcg	15		101	
Selenium, mcg	2		14	
Sodium, mg	30		203	
Sodium, mEq	1.3		8.8	
Potassium, mg	107		723	
Potassium, mEq	2.7		18.2	
Chloride, mg	65		439	
Chloride, mEq	1.8		12.2	

For more information, contact your Abbott Nutrition Representative or visit
www.abbottnutrition.com

Preparation

Instructions for Use:

Your baby's health depends on carefully following these directions. Failure to follow these directions could result in severe harm. Ask your baby's doctor if you need to use cooled, boiled water for mixing and need to boil (sterilize) bottles, nipples and rings before use.

Concentrated Liquid

- 1 Rinse can lid; shake very well; open with a clean punch-type opener.
- 1 Pour equal amounts of formula and water into bottle.
- 1 Attach nipple; swirl gently to mix. Once feeding begins, use within 1 hour or discard.

Ready to Feed

- 1 **Do not add water.**
- 1 **Do not use if band around cap or inner foil seal is damaged.**
 - 1 Shake very well before each use. Remove protective band, twist off and clean cap.
 - 1 Invert cap; press down to pierce foil, then turn cap a half turn. Remove foil.
 - 1 Pour formula into bottle; attach clean nipple. Once feeding begins, use within 1 hour or discard.
- 1 Prepare and feed all formula from 1 QT bottle within 3 days after opening.

Storage Instructions:

Concentrated Liquid

- 1 Cover opened cans and prepared formula not used immediately.
- 1 Store in refrigerator and **feed to baby within 48 hours.**
- 1 Store unopened cans at room temperature; avoid extreme temperatures.

Ready to Feed

- 1 Once opened, store quart bottle in refrigerator.
- 1 Store prepared bottles in refrigerator and **feed to baby within 48 hours.**
- 1 Store unopened at room temperature; avoid extreme temperatures.
- 1 Do not reuse bottle.