Infant formula with iron

Product Information: Similac Sensitive® Powder

Infant formula with iron

- SIMILAC SENSITIVE is a 20 Cal/fl oz, nutritionally complete infant formula that is an alternative to standard milk-based formulas. For fussiness, agas, or mild spit-up. Suitable for infants with lactose sensitivity.
- Easy-to-digest gentle carbohydrate for fussiness and gas due to lactose sensitivity.
- OptiGRO® is our exclusive blend of DHA, lutein and vitamin E: these important ingredients are found in breast milk.
 - O DHA for brain and eye development.
 - O Lutein to support eye health.
 - Vitamin E to support developing cells.
- 1 Our exclusive formula has:
 - O Calcium for strong bones no palm olein oil.
 - O Nucleotides to help support the immune system.
 - O Prebiotics to help promote digestive health.
 - O Carotenoids like those naturally found in breast milk.
- 1 Clinically shown to support normal growth in infants.¹
- 1 A unique blend of carbohydrates using two different absorptive pathways to help maximize absorption and minimize malabsorption risks.
- Low osmolality (200 mOsm/kg water).
- 1 Halal.
- 1 Kosher; made with dairy ingredients.

Safety Precautions

- 1 Not for infants or children with galactosemia.
- 1 Never use a microwave oven to warm formula. Serious burns can result.
- Powdered infant formulas are not sterile and should not be fed to premature infants or infants who might have immune problems unless directed and supervised by your baby's doctor.

Ingredients

Powder Unflavored:

Corn Syrup, Milk Protein Isolate, High Oleic Safflower Oil, Sugar, Soy Oil, Coconut Oil. Less than 2% of: Schizochytrium Sp. Oil*, M. Alpina Oil†, Short-chain Fructooligosaccharides‡, Beta-Carotene, Lutein, Potassium Chloride, Sodium Citrate, Calcium Phosphate, Potassium Citrate, Soy Lecithin, Magnesium Phosphate, Choline Chloride, Ascorbic Acid, Inositol, Ascorbyl Palmitate, Ferrous Sulfate, Taurine, Calcium Carbonate, Choline Bitartrate, Zinc Sulfate, L-Carnitine, Mixed Tocopherols, Niacinamide, d-Alpha-Tocopheryl Acetate, Calcium Pantothenate, Magnesium Chloride, Vitamin A Palmitate, Copper Sulfate, Thiamine Hydrochloride, Riboflavin, Pyridoxine Hydrochloride, Folic Acid, Potassium Iodide, Manganese Sulfate, Phylloquinone, Biotin, Sodium Selenate, Vitamin D3, Vitamin B12, Potassium Hydroxide, and Nucleotides (Adenosine 5'-Monophosphate, Cytidine 5'-Monophosphate, Disodium Guanosine 5'-Monophosphate, Disodium Uridine 5'-Monophosphate).

Allergens: Contains milk ingredients.





^{*} Due to lactose sensitivity.

 $^{^\}dagger$ Typical value 0.1 g lactose/100 kcal compared to Similac® Advance® (11 g lactose/100 kcal).

¹ Lasekan JB, et al. Clin Pediatr. 2011;50(4):330-337.

Source of DHA.

[†] Source of ARA.

[‡] Prebiotic.

Infant formula with iron

Availability

List Number	Item
57539	Similac Sensitive Powder / 12.5 oz (354 g) can / 6 ct
62951	Similac Sensitive Powder / 29.8 oz (845 g) can / 6 ct
68082	Similac Sensitive Powder / 20.1 oz (570 g) Container / 4 ct



Infant formula with iron

Nutrition Information - Powder Unflavored

Nutrition Information - Powder Unflavored						
	100 Cal (5 fl oz, pr	epared as directed)	1000 mL (1.1 QT, prepared as directed)			
	Value	%RDI / %DV	Value	%RDI / %DV		
Calories	100		676			
Volume, mL	148		1000			
Protein, g	2.1		14.19			
Fat, g	5.4		36.49			
Carbohydrate, g	10.9		73.65			
Water, g	133		899			
Linoleic Acid, mg	1000		6757			
Vitamin A, IU	300		2027			
Vitamin D, IU	60		405			
Vitamin E, IU	1.5		10.1			
Vitamin K, mcg	8		54.1			
Thiamin (Vitamin B1), mcg	100		676			
Riboflavin (Vitamin B2), mcg	150		1014			
Vitamin B6, mcg	60		405			
Vitamin B12, mcg	0.25		1.69			
Niacin, mcg	1050		7095			
Folic Acid (Folacin), mcg	15		101.4			
Pantothenic Acid, mcg	450		3041			
Biotin, mcg	4.4		29.7			
Vitamin C (Ascorbic Acid), mg	9		60.8			
Choline, mg	24		162.2			
Inositol, mg	24		162.2			
Calcium, mg	84		568			
Calcium, mEq	4.2		28.4			
Phosphorus, mg	56		378			
Magnesium, mg	6		40.5			
Iron, mg	1.8		12.16			
Zinc, mg	0.75		5.07			
Manganese, mcg	5		33.8			
Copper, mcg	90		608			
lodine, mcg	15		101			
Selenium, mcg	2		14			
Sodium, mg	30		203			
Sodium, mEq	1.3		8.8			
Potassium, mg	107		723			
Potassium, mEq	2.7		18.5			
Chloride, mg	65		439			
Chloride, mEq	1.8		12.2			

For more information, contact your Abbott Nutrition Representative or visit www.abbottnutrition.com



Infant formula with iron

Preparation

Instructions for Use:

Your baby's health depends on carefully following these directions. Proper hygiene, handling and storage are important when preparing infant formula. Failure to follow these directions could result in severe harm. Ask your baby's doctor if you need to use cooled, boiled water for mixing and if you need to boil (sterilize) utensils, bottles, nipples and rings before use.

- 1 Wash your hands, surfaces and utensils. Pour water into clean bottle (see mixing guide).
- 1 Add 1 unpacked level scoop (8.8 g) to each 2 fl oz of water. Return dry scoop to can.
- 1 Cap bottle; shake well; attach clean nipple.
- 1 Once feeding begins, use within 1 hour or discard.

Powder Mixing Guide

Measure water	+	Add scoop(s) of <i>unpacked level powder</i> using enclosed scoop
2 fl oz		1 scoop (8.8 g)
4 fl oz		2 scoops
6 fl oz		3 scoops
8 fl oz		4 scoops

Each scoop adds about 0.2 fl oz to the amount of prepared formula. For larger size mixing instructions, please visit www.Similac.com/mixing info.

When mixed as directed:

- 12.5 oz container makes approximately 90 fl oz of formula.
- 20.1 oz container makes approximately 145 fl oz of formula.
- 29.8 oz container makes approximately 215 fl oz of formula.

STORAGE:

- Once mixed, store bottles in the refrigerator; feed to baby within 24 hours.
- 1 Store unopened or opened container at room temperature; avoid extreme temperatures.
- Use opened can contents within 1 month.

