

СПИСОК ИССЛЕДОВАНИЙ SIMILAS:

1. **К. Геринг 2016:** Goehring, K. C., B. J. Marriage, J. S. Oliver, J.A. Wilder, E. G. Barrett and R. H. Buck (2016). "Similar to Those Who Are Breastfed, Infants Fed a Formula Containing 2'-Fucosyllactose Have Lower Inflammatory Cytokines in a Randomized Controlled Trial." J Nutr 146(12): 2559- 2566.
2. **Б.Мэрридж с соавт. 2015 г.:** Marriage, B. J., R. H. Buck, K. C. Goehring, J.S. Oliver and J. A. Williams (2015). "Infants Fed a Lower Calorie Formula With 2'FL Show Growth and 2'FL Uptake Like Breast- Fed Infants." J Pediatr Gastroenterol Nutr 61(6): 649-658.
3. **Э.Реверри 2018:** Reverri EJ, Devitt AA, Kajzer J, Baggs G, Borschel MW. Review of the Clinical Experiences of Feeding Infants Formula Containing the Human Milk Oligosaccharide 2'-Fucosyllactose. Nutrients 2018, 10, 1346; doi:10.3390/nu10101346
4. **Б.Лойд с соавт. 1999:** Lloyd B, Halter RJ, Kuchan MJ, et al. Formula tolerance in post breastfed and exclusively formula-fed infants. Pediatrics. 1999;103:e7
5. Data on File, AK70. Abbott Nutrition, Abbott Laboratories, 2017